SARS-CoV-2 (2019-nCoV) vaccine, Gen 13

Rapid Deployment Vaccine Collaborative (RaDVaC)

Preston Estep, Don Wang, Alex Hoekstra, Ranjan Ahuja, George Church, Brian Delaney, and over 100 others

Table of Contents

TERMS OF USE (updated 2022-07-21) Overall RaDVaC project goal (updated 2022-05-09) Vaccines and the COVID-19 global emergency (updated 2022-05-12) VACCINE BACKGROUND AND DESIGN (updated 2022-05-16) POSSIBLE MECHANISMS OF VACCINE-ENHANCED DISEASE, VACCINE-INDUCED AUTOIMMUNITY, AND MITIGATION STRATEGIES (updated 2020-10-24) KEY TECHNICAL FEATURES AND SPECIFICATIONS (Updated 2021-03-06) GOALS (last updated 2022-01-14) MATERIALS AND METHODS (updated 2021-03-07) Selection and purchase of the vaccine ingredients Tools for designing and analyzing vaccine components (NEW 2022-07-22) Preferred equipment and materials (updated 2020-11-04) Selection of epitopes (updated 2021-03-17) Peptides and vaccine composition (last updated 2022-01-14) Peptide synthesis general guidelines Gen 10 and forward (updated 2022-01-14) Immunogenic enhancements Gen 13 updates (updated 2022-08-13) Gen 13 Core peptides Gen 13 Optional peptides Spike or S protein amino acid sequences represented by epitope peptides Wuhan parental strain **Omicron variants** Nucleocapsid protein epitopes **T-cell epitope peptides** SUMMARY OF THE PROTECTIVE STRATEGY OF THE GEN 13 VACCINE DESIGN (updated 2022-07-20) PREPARATION OF CHITOSAN NANOPARTICLE VACCINE (updated 2022-07-22; update includes note about the adjuvant imiquimod) ADMINISTRATION OF THE VACCINE (updated 2021-03-07) ASSESSMENTS OF IMMUNE RESPONSE (updated 2021-03-07) Samples to be collected and tested Antibodies and B-cell immune response **B-cell epitopes and ELISA antigen targets** Nucleocapsid protein B-cell epitopes / antigen targets **T-cell immune response** SUPPLEMENTAL MATERIALS **ADDITIONAL SEQUENCES** Wuhan strain Spike protein sequence Nucleocapsid protein sequence

TERMS OF USE (updated 2022-07-21)

This document describes the rationale, design, formulation, and possible self-administration of an experimental vaccine candidate for SARS-CoV-2. By using this information you agree to the following: 1) you are a consenting adult (in the USA, at least 18 years of age); and 2) you take full responsibility for your use of RaDVaC information and vaccine or material--including redistribution, modification, vaccine formulation, production and administration.

A foundational principle of the vaccine development and deployment strategy of RaDVaC is rapid iteration and testing of vaccine designs, based on newly published information from the forefront of biomedical research. This agile approach has the potential to produce better vaccines much more quickly than traditional approaches. However, such rapid design improvements currently are not compatible with established clinical trial requirements of fixed vaccine design, nor with review and approval by ethics committees (due to shifting vaccine design and continually updating self-experimentation and testing protocols). Therefore, the information presented within has not been approved by an institutional review board or any other type of ethics committee, and you understand and agree that any implementation of this information constitutes self-experimentation.

The purpose of this open-source vaccine effort is to reduce risk of harm from SARS-CoV-2. In addition to providing the results of our own research and experimentation, we hope to motivate others to build on our work, and to pursue diverse evidence-based approaches. Given the immense complexity and variability of individual human biology, it is not possible to predict all potential physiological responses to any vaccine. But as is true for most or all vaccines used on a large scale, there is a tradeoff between a larger known risk from the disease the vaccine is designed to prevent or mitigate, and the smaller risk introduced by the vaccine itself. Any vaccine poses risks, and, if used in enough people, will cause some degree of harm. Furthermore, certain harm, such as allergic and possibly anaphylactic response, will be readily seen and measured, whereas benefit is more difficult and takes longer to assess. These vaccine candidates are no different in this regard; and because quality of delivery is highly dependent on the meticulousness of individual end users, the vaccines might pose unique risks not posed by typical commercial vaccines.

Does Not Constitute or Substitute for Medical Advice

Information presented here is ongoing research, and is not intended as a substitute for medical advice. RaDVaC is not responsible for the decision to administer, or to receive administration of, any vaccine.

No Promises or Guarantees of Efficacy

Vaccines are often received with the false hope of efficacy, without testing to determine the degree of individual immune response. For example, influenza killed about 100,000 people in the U.S. between late 2016 and early 2018. Yet, the influenza vaccines available in that period were substantially less than 50% effective against H3N2, the flu strain mainly responsible for the death toll. Many who died were vaccinated but not protected from the virus, and testing for vaccine-induced immunity was essentially non-existent. Vaccine-induced immunity can be more challenging to assess than immunity due to viral infection, and such is probably the case for the nasal vaccine described here. Because this work is a research undertaking, no expectation is given regarding any degree of efficacy in granting protection against SARS-CoV-2. On the contrary, see possible risks and uncertain benefits below.

Preventive, Not Therapeutic

Even if this vaccine works as intended, it will not help someone who has been infected. It will only work as a preventive measure taken weeks in advance of virus exposure.

Not a Clinical Trial

Any use of the information in this document, or provided in correspondence with us does not constitute 'partnership' or 'recruitment' for an organized trial in any way. The supported conjecture here is given as a starting point for additional individual or organized, sponsored research. As of public release of this information, no organized clinical trials have been performed to test this vaccine.

Not Approved or Reviewed by the FDA

The information, procedures, and conclusions presented here have not been approved, or reviewed by the FDA, or any other regulatory body.

Possible Risks and Uncertain Benefits

- Immediate allergic or other serious reaction
- Unforeseen long-term effects
- Instillation/administration of the vaccine in an inappropriate way or in an infected area might increase the risk of infection by enhancing viral entry into the body
- Benefits are uncertain. There are extensive published histories of the materials and procedures described in this document, but every novel vaccine should be considered experimental, with the possibility there will be no benefit.

- Even if there are signs of immune response there is no guarantee this response is indicative of protection from SARS-CoV-2 infection, or if protection is achieved, how long it will last.
- Even if the vaccine confers protection from the virus, certain methods for assessing protection--such as measuring antibodies due to previous infection--might not capture vaccine-induced immunity. In such cases, infection, and thus a positive test result, are unlikely to occur. Immunity passports and other privileges given to convalescents might be difficult to obtain for those with vaccine-induced immunity.
- Use of this vaccine candidate may change the efficacy of any future vaccines you may take or that are administered to you, in unknown ways

No Offer of Service and No Access to Materials

All information in this document or via correspondence is made available as open science. We do not sell equipment or services, e.g. we will not prepare this formulation or any other on behalf of a requestor. All vaccine materials at our disposal are kept in a secured laboratory. We cannot and will not keep materials outside the lab for easy access or distribution. We will not provide laboratory or equipment access for vaccine production.

Probability and Coincidence

Vaccines have become associated with negative health events that do not result from the vaccine, but that happen to occur soon after administration. Negative health events will occur with baseline frequency even in the absence of vaccination. Increasing use of any vaccine will increase the probability of vaccine recipients experiencing a negative health event purely by chance, including shortly after vaccination. By using the information presented here, you acknowledge the increasing likelihood of such coincidences, and assume full responsibility for any use of the information herein and for real or perceived negative events, irrespective of the cause.

Self-Experimentation

By utilizing the information and scientific opinions in this document, and in any correspondence with us, you acknowledge and agree that any use to develop and self-administer a substance is an act of self-experimentation. Laws and regulations on such actions may differ in your national or state jurisdiction, and it is your responsibility to be informed about them.

Additionally, the type of work described in this document requires certain equipment and level of skill with laboratory techniques. You agree to assume full responsibility for acquiring proper equipment, knowledge, and training, and for attempting to formulate or administer vaccine.

Licenses

All contents of the RaDVaC website and white papers are covered by a Creative Commons Attribution 4.0 License (CC BY 4.0), and by an Open COVID License – Patent 1.1 (OCL-P v1.1). You are free to Share (copy and redistribute the material in any medium or format), and Adapt (fork, alter, and build upon) the material for any purpose. The only requirement is Attribution.

Overall RaDVaC project goal (updated 2022-05-09)

The Rapid Deployment Vaccine Collaborative (RaDVaC) was founded by members of the Mind First Foundation in March of 2020 in order to address the outbreak of the SARS-CoV-2 virus, which quickly became a global emergency. The SARS-CoV-2 pandemic produced many notable accomplishments, including the most rapid large-scale vaccine trialing and emergency deployments in history. Nevertheless, vaccines were only available to selected members of the public in high-income countries about a year after initial signs of the emerging pandemic. This year-long "vaccine access gap" allowed SARS-CoV-2 to spread largely unchecked throughout the world.

The goal of RaDVaC is to create a modular vaccine platform for extremely rapid deployment at the beginning of a serious outbreak of any pathogen for which no good vaccine is available. The foundational philosophy of RaDVaC is that the best way to accomplish this goal is to freely, openly, and publicly share all of its work on vaccines and trialing; and to advance a distributed, global network of researchers engaged in agile R&D. For many individual researchers, public health authorities, and others involved in vaccine development and deployment, this will include localized and decentralized vaccine production and trial design. While the SARS-CoV-2 pandemic was the catalyst for the initiation of the RaDVaC project, it will not be the last serious outbreak; it is critical to establish a reliable scientific and technological foundation for rapid, decentralized vaccine deployment in future scenarios. This long-term endeavor will necessitate paradigm shifts in vaccine R&D, testing, and production, as well as in data sharing and regulatory frameworks.

We intend for our work to comprise an evolving open-source toolkit (or Vaccine Developer Kit / VDK) to enable vaccine R&D, production, testing, approval, and deployment that is faster, less expensive, and more adaptable and agile to viral variants and regional needs. As proponents of citizen-science and deeper public engagement in scientific issues, we envision a scientific and medical culture that promotes greater access and participation, and which is more scientifically robust and resilient than the current norm. Ultimately, we seek to model and promulgate open-source practices of information- and expertise-sharing so that global challenges of all kinds can be addressed more effectively.

Vaccines and the COVID-19 global emergency (updated 2022-05-12)

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2; a.k.a. 2019-nCoV; disease: COVID-19) is responsible for a worldwide pandemic far beyond the scope of any other public health crisis in over a century. As of May, 2022, the official worldwide COVID-19 death toll is over 6 million. According to analyses by the Institute for Health Metrics and Evaluation, and separately,

by Samira Asma, Assistant Director-General of the WHO data and analytics division, the true number of deaths caused by COVID-19 is likely more than three times the official count^{1,2}.

Higher-income countries now have excellent vaccine access, although substantial fractions of populations in almost all countries refuse to vaccinate. Vaccine access is far worse in low- and middle-income countries. These countries had only minimal access to commercial vaccines through 2021, and will likely not have sufficient access to these vaccines until well after 2022³. These significant delays not only lead to continued deaths and chronic illness, but also allow additional viral strains, including escape mutants to arise, compounding the scientific and public health challenges for both low-income and high-income nations, and threatening the fragile progress toward recovery as these new variants spread across the globe.

Because of these large, geographically distributed reservoirs, experts are forecasting that SARS-CoV-2 is transitioning toward being globally endemic. New variants are predicted to emerge regularly for the foreseeable future, causing case and mortality spikes similarly to influenza, though with higher absolute mortality. Only effective, broad-spectrum coronavirus vaccines have the potential to bring this deadly transition phase to an end. The unvaccinated might better benefit from such next-generation vaccines targeting newer variants, rather than settling for inferior protection from vaccines designed for variants no longer in circulation.

In a pandemic situation, ideally the first choice one must make in the decision to use a vaccine is which of the two possible ways one will first be exposed to immunogenic elements of a pathogen: by infection, or by vaccination. Vaccines are designed to stimulate an immune response that provides long-lasting protection against infection and/or disease by a pathogen. While infection by that pathogen and recovery from disease can sometimes also bestow such protection, pathogens, including viruses, possess many complex mechanisms for hijacking cellular and physiological functions and evading immune responses. Immunity gained from pathogen infection can therefore be suboptimal, or even dysfunctional due to these pathogen tactics. The same mechanisms can compromise health, and cause prolonged suffering and possibly death. We believe therefore that vaccine immunity is a far better choice; but in the current pandemic that option was available to almost no one for more than a year, and is still not available to much of the global population.

There is substantial published information on all aspects of vaccine production, testing, and delivery, an increasing amount of which is specific to the SARS-CoV-2 virus. In addition, thanks to generous grants from multiple organizations, including the Bakewell Foundation and the Balvi

¹ <u>https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)02796-3/fulltext</u>

² Samira Asma, WHO press briefing, 2021-05-21. <u>https://www.reuters.com/article/idUSL5N2N81LY</u>

³ <u>https://www.theguardian.com/society/2021/jan/27/most-poor-nations-will-take-until-2024-[...]-immunisation</u>

Filantropic Fund,⁴ we will be able to extensively assess new versions of the vaccine with preclinical safety and immuno-efficacy studies of our own.

Vaccines are the safest of all therapeutic classes. Because of their safety and efficacy, they have by far the highest probability of success in the overall course of clinical trials (over 40%, which is about twice the rate of the next highest class).⁵ Because of this and the impact of the pandemic, commercial trials have been greatly accelerated.

Nevertheless, commercial vaccines must be designed to enable large scale production and deployment, and they require regulatory approval for sale, both of which greatly constrain and slow progress. Critically, certain features of commercial vaccines aren't required for research-level production and testing; therefore, a much shorter timeline is possible for a smaller scale, self-administered vaccine. Herein we describe the formulation of a vaccine using proven and mostly off-the-shelf components, with a low per-dose cost.

This will be a versioned living document, embodying and enabling incremental improvements in vaccine design and testing.

VACCINE BACKGROUND AND DESIGN (updated 2022-05-16)

There are many advances in vaccine research and technology development that have not appeared in a commercial product, in part because commercial vaccine design and production is constrained by different factors than a small-scale research vaccine. Desirable features of a RaDVaC open-source vaccine are high safety and efficacy, low cost, and ease of production and administration. A key feature of the RaDVaC vaccine is rapid iteration in order to update designs as new discoveries about the biology and clinical manifestations of SARS-CoV-2 are published in the biomedical literature and on preprint servers. This feature is critical to the RaDVaC mission of rapid development and deployment of effective vaccines. We address all of these features in this document, and consider subcategories, such as near-term and long-term safety.

Key issues and some differentiating factors between commercial vaccines and the RaDVaC research vaccine platform:

• **Rapidly responsive design.** A key feature of our rapid deployment strategy is the flexibility of the vaccine platform, allowing very fast and efficient updating of vaccine designs in response to recent research. On multiple occasions we have updated vaccine

⁴ <u>https://radvac.org/wp-content/uploads/2022/05/Balvi_RaDVaC-announcement.pdf</u>

⁵ <u>https://projectalpha.mit.edu/pos/</u>

designs based on important information published in the biomedical literature only weeks or days earlier.

One example is the important discovery showing that T-cell responses are found in patients almost two decades after recovery from SARS-CoV-1 (a very closely related coronavirus)⁶. The portions of the virus responsible for these very long-lived T-cell responses (CD4 and CD8) are in highly conserved parts of the virus. They are not present in the Spike protein, which is not highly conserved, and which is the primary target of leading vaccines in the current race to commercialization. A second publication reveals the identities of 29 experimentally verified, immunodominant CD8 T-cell epitopes of SARS-CoV-2⁷. This publication confirms several important findings of a prior preprint⁸, and some of these epitopes are shared among the small group of long-lived epitopes from SARS-CoV-1, none of which resides in the Spike protein. Only 3 of 29 epitopes lie within the Spike protein, and only one lies within the RBD region of Spike. Another subsequent publication experimentally confirmed and extended these findings, also showing that Spike epitopes are not among the top 10 immunodominant CD8 T-cell epitopes⁹. These findings suggest that the Spike protein might not provide extensive and robust T-cell protection, and that RBD will provide even less protection for many people, especially the large numbers of people with MHC haplotypes not represented by these antigens.

Consistent with this expectation, data on the leading commercial vaccines suggest that they do not confer sterilizing immunity (the immune system's ability to stop a pathogen from entering the body, or if it gains entry, from replicating within the body). Again, the RaDVaC strategy allows for frequent updating of vaccine designs based on ongoing research throughout the world, so that the level of protection keeps pace with new knowledge, and new variants.

Near-term safety. Near-term safety of the intranasal vaccine described here should be excellent when prepared appropriately. Intranasal vaccines have a long history of demonstrated safety. The vaccine formulations described here can also be used for other delivery routes, including inhalation, oral, or parenteral delivery (injection), but these, especially the latter, should be attempted only by skilled practitioners. In general, a vaccine is more safely used for inhalation into the lungs¹⁰, oral, or intranasal delivery, than for injection. This document is focused on intranasal delivery, but experts will know how

⁶ https://www.nature.com/articles/s41586-020-2550-z

⁷ <u>https://www.medrxiv.org/content/10.1101/2020.07.24.20161653v2.article-info</u>

⁸ https://www.biorxiv.org/content/10.1101/2020.06.05.134551v1.full.pdf

⁹ https://www.medrxiv.org/content/10.1101/2020.07.31.20165647v2.article-info

¹⁰ <u>https://www.sciencedirect.com/science/article/pii/S1549963415000313</u>

to adapt the information herein to another delivery route. For any delivery mode, sourcing of high-quality materials and meticulous preparation are essential for maintaining vaccine safety.

- Long-term safety. The long-term safety of any new vaccine is currently difficult or impossible to predict. Even widely deployed commercial vaccines have resulted in serious and unforeseen complications. Vaccines that have shown serious side effects are injectable whole virus or subunit formulations. There are at least three potentially serious complications that might arise in the long term: tolerance, vaccine-enhanced disease (VED), and adjuvant-triggered immune or neurological complications.
 - Immune tolerance denotes diminished immunity resulting from exposure to an antigen. This attenuated immune response is commonly observed in food antigens, in "self" antigens of one's own body, and in antigens from commensal organisms. In general, it is thought that extremely large and/or frequent exposures and oral doses lead to tolerance. While in the above examples, immune tolerance is desirable, in the case of vaccines it can cause the immune system not to mount an appropriate defensive response to infection.
 - Vaccine-enhanced disease (VED). A small number of injected vaccines have led to enhancement of disease, meaning that infectivity is enhanced, or the disease is made more serious in people who have been vaccinated, relative to unvaccinated controls. Possible mechanisms underlying VED, such as antibody-dependent enhancement (ADE) are explored at length in the section below, together with various vaccine design features and strategies for reducing the likelihood of VED.
 - Adjuvant hyperstimulation or toxicity. Adjuvants help stimulate a robust immune response to a vaccine; however, certain adjuvants have caused hyperstimulation and other serious side effects. For example, alum produces a robust Th2 immune response, but an unbalanced ratio of Th2:Th1. A Th2 polarized response, and alum in particular, have been implicated in immunopathology, including ADE. Adjuvants can also be toxic. As one example, the intranasal use of a detoxified mutant form of *Escherichia coli* Heat Labile Toxin has resulted in transient Bell's palsy, or facial nerve paralysis.¹¹ Specific adjuvants have also been suggested to play a role in the development of autoimmunity. For example, the influenza vaccine Pandemrix appears to have increased the incidence of narcolepsy in Scandinavian countries ¹². The leading explanation is that the vaccine triggered a CD4-mediated autoimmune reaction against sleep-regulating neurons (although the vaccine-attributable risk is very low, with 1 case in 18,400 vaccinations, or about

¹¹ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2737308/</u>

¹² https://pubmed.ncbi.nlm.nih.gov/28847694/

0.005%). One suggested contributing factor is the strong squalene-based adjuvant used in Pandemrix. The primary reason for using strong adjuvants in commercial vaccines is to avoid the need for boosters, by triggering a robust immune response with a single dose.

- **Stability.** Stability is a key determinant of a commercial vaccine. Formulations that are both safe and effective in a research setting, but with limited shelf life, are generally excluded from commercial products. We have discovered that there are formulations for vaccines that are extremely simple, safe, and effective, but have only short-term stability (on the order of weeks). For example, chitosan gel nanoparticles have been shown to be effective and extremely simple to formulate, but short shelf life has contributed to their limited use in commercial vaccines.
- Ease of delivery. Intranasal vaccines have advantages over other delivery approaches, including the most common delivery modality, parenteral/injection. Intranasal delivery has been demonstrated to be very safe, with mild side effects typically equal to those seen in placebo-treated subjects. Importantly, it can elicit not only systemic immunity, but also mucosal immunity at the point of infection for respiratory viruses.¹³ Commercial intranasal influenza vaccine is available, and relative to parenteral (injected) administration shows equal efficacy for systemic immunity, but greater efficacy for mucosal immunity at sites of entry (nose, lungs) for respiratory viruses.¹⁴ This is critically important for SARS-CoV-2 because early studies show that a majority of infections begin in the nasal cavity¹⁵. As of May 2022, most commercial SARS-CoV-2 vaccines in development are designed to be injected (parenteral), a route which is unlikely to provide mucosal immunity to infection. As of early 2022, we have also been exploring other methods of delivery that we expect to be easy to administer, safe, and effective, including sublingual, oral, inhaled, and transdermal delivery systems.
- Prime-Boost. Alternative methods of delivery can be as effective as injection, but to achieve the same level of immune response and protection--especially with milder vaccines and adjuvants--multiple doses are often required. The initial dose is the "prime" and subsequent doses are given to "boost" or increase the immune response. Prior exposure to a pathogen or closely related one has a similar effect to a prime dose of vaccine. The only commonly used intranasal vaccine is for influenza, and since essentially everyone is naturally exposed to influenza, the nasal vaccination with attenuated virus is in essence a booster. Necessity of multiple boosters has limited the commercial production of intranasal (and other mildly/moderately immunostimulating) vaccines for

¹³ https://pdfs.semanticscholar.org/6e55/db027b08b367eaac55ec54a730f4c99061ec.pdf

¹⁴ https://www.tandfonline.com/doi/abs/10.1586/erv.12.31

¹⁵ <u>https://www.cell.com/cell/pdf/S0092-8674(20)30675-9.pdf</u>

which prior ubiquitous exposure is unlikely. An important corollary to a multiple-dose design is titration of adequate immune response for each individual. Whereas the immunogenicity-safety profile of commercial formulations must be standardized to an "average" level--overdosing some and underdosing others--with accompanying testing the RaDVaC approach should allow for more precise dosing.

• Efficacy. Intranasally delivered vaccines have shown mild side effects and high levels of efficacy of both mucosal and systemic immunity, when delivered in a prime-boost regimen (in both animal models and human trials; see one review by Smith et al.¹⁶). One of our research priorities is to determine when other non-parenteral/injection delivery modes will prove to be equally safe and efficacious.

Given the above key points, it is clear that a rapid-deployment vaccine that is both safe and effective is likely to have multiple attributes that differentiate it from a commercial vaccine. And it becomes clear why the formulation described here has not been used in a commercial product; it is primarily not lack of safety or efficacy, but other factors related to stability, scalability, patient compliance, profitability, and regulatory complexities of commercial vaccines. Commercial vaccines are designed to produce high levels of immunity with a single dose. A prime and multi-boost schedule is used routinely in the research setting, but concerns about compliance (failure to boost) limit commercial deployment.

For multiple reasons, including those given above, non-parenteral/non-injection delivery is difficult to achieve for a commercial vaccine. However in the research context it has shown multiple advantages. To start, relative to injection, there is no risk of needle injury or transmission of blood-borne infection. Further, it is not only relatively easier to administer, but in certain cases and for some pathogen classes like respiratory viruses, intranasal or other mucosal delivery modes are in fact the preferred methods of delivery. Immunization by the intranasal route, for example, not only prevents viral infection through the nasal membranes, but also efficiently stimulates a strong mucosal immune response in the upper respiratory tract and lungs. Sublingual and other mucosal delivery modes are also likely to induce a strong mucosal immune response. As one example, Gai and colleagues have shown that a SARS vaccine delivered intranasally elicits a robust mucosal immune response, protecting against initial infection, whereas the same vaccine delivered by injection does not.¹⁷ This difference is important because the area of the mucosal surfaces (nasal, lungs, gastrointestinal, urogenital, etc.) is very large, about 200 times the surface area of the skin, and about 70% of pathogens enter through these routes.¹⁸ Compliance is also very high because intranasal delivery doesn't involve needles or cause pain or significant

¹⁶ https://www.tandfonline.com/doi/full/10.4161/hv.27449

¹⁷ <u>https://scholar.google.com/scholar?cluster=13722847195977601349&hl=en&as_sdt=0,22</u>

¹⁸ https://pdfs.semanticscholar.org/6e55/db027b08b367eaac55ec54a730f4c99061ec.pdf

discomfort. High safety and ease of administration are expected to contribute to high rates of immunization. For a recent review of research on nasal nanovaccines, see Bernocchi et al¹⁹ (Table 1).

Synthetic peptide-based vaccines permit a more curated and intentional immunogen design than do vaccine platforms which are based on attenuated whole viruses or even full-length Open Reading Frames (ORFs) of key epitope proteins. In commercial vaccines, there is a trend toward highly defined, minimal designs. According to Pompano and colleagues "[v]accine design is moving away from using whole pathogens in favor of selecting only the most protective antigens for immunization with a suitable adjuvant. The most tailored of these sub-unit vaccines utilize specific B-cell or CD8+ T-cell epitopes, such as the short peptides and carbohydrates that have been used to elicit antibody or killing responses against malaria pathogens, bacterial infections, or tumors. In addition to the desired target epitope, these vaccines also require one or more CD4+ T-cell epitopes to engage CD4+ helper T-cells."²⁰

Such compact and efficient designs are not achieved easily, and require substantial data on the immunogenicity of all epitopes of a pathogen, but such designs have the advantage of limiting pathogenic effects of certain viral antigens. For instance, coronavirus Spike and Nucleocapsid full-length proteins have been associated with ADE in animals and human cellular studies. Yasui and colleagues showed that vaccination with Nucleocapsid protein does not provide protective immunity, yet enhances immunopathology.²¹ Vaccination with certain epitopes of Spike protein does confer protection, but use of the full-length Spike might be problematic. For example, from Tais and colleagues: "...full-length S protein should be used with caution. Kam et al. reported that although a recombinant, trimeric SARS-CoV S protein vaccine elicited a protective immune response in mice the anti-S antibodies also mediated antibody-dependent enhancement of viral entry into human B-cells in vitro. In another study, ferrets vaccinated with SARS-CoV full-length S protein expressed by a recombinant modified vaccinia Ankara grown in BHK21 and Vero E6 cells show enhanced virulence of hepatitis induced by SARS-CoV. Furthermore, the use of a SARS S protein vaccine may lead to enhanced disease and immunopathology instead of protection as seen for feline coronavirus, feline infectious peritonitis virus. Given these concerns, the use of a SARS vaccine strategy in which the full-length S protein is used may not be optimal for humans. Hence, the best approach would probably be to use small S protein epitopes that are major neutralization determinants."22 Vaccines against other pathogens that similarly use full-length constructs have also elicited enhancement of viral sensitivity or disease. Nevertheless, as of mid

¹⁹ <u>https://www.ncbi.nlm.nih.gov/pubmed/28698066</u>

²⁰ https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4227912/

²¹ https://www.jimmunol.org/content/181/9/6337

²² https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2756483/

2022, billions of people have been vaccinated with full-length Spike vaccines, and the frequency of serious side effects is minimal. Furthermore, as of Dec 2021, the combined seroprevalence of vaccination and/or infection reached about 95% of the adult population in the US.²³ Global exposure in 2022 approaches a similarly high frequency. Therefore, exposure to full-length Spike is the norm, and one can either be exposed through vaccination or through infection. We believe it is vastly preferable to be vaccinated than infected. The vaccine design described here is based on key epitopes in the full-length Spike protein, in addition to predicted and experimentally validated T-cell epitopes. If you decide to modify the design detailed herein, including the use of modified or full-length Spike protein as an immunogen, please let us know and keep us informed of additional salient information.

Synthetic peptide synthesis provides freedom to design epitopes of sufficient length for immunogenic stimulation, but predicted not to trigger these serious side effects. This approach also allows the use of multiple epitope peptides, either concatemerized in a single, multimeric linear peptide, or as a collection of individual peptides. The approach taken here uses B-cell epitope peptides of S protein sequences, and T-cell epitopes from S and other proteins. Most or all of the B-cell epitope amino acid sequences contain predicted T-cell epitopes. We have employed many of these combination epitopes in order to keep the design compact and relatively simple and inexpensive to produce. Later versions of epitopes were selected based on data for neutralizing antibodies and T-cell responses from previously SARS-CoV-2 infected convalescent patients.

Immunopathology isn't exclusively determined by immunogen, however. It can also be influenced greatly by adjuvant choice. Antibody-dependent enhancement of disease and immunopathology appears to be mediated primarily by a Th2 (or Th17) response, and opposed by a Th1 response ²⁴. Extensive research on Respiratory Syncytial Virus, SARS, and other respiratory viruses shows that ADE immunopathology will be more likely with the use of alum or other Th2-boosting adjuvant.²⁵ From Honda-Okubo et al: although an anti-SARS vaccine "formulated with alum protected against mortality, these mice developed severe lung eosinophilia at day 6 post-challenge, reminiscent of the lung pathology induced by alum-adjuvanted RSV vaccines and indicating alum's general lack of suitability as a coronavirus vaccine adjuvant." These concerns make alum alone a poor adjuvant choice. Yet alum is not without substantial advantages. Alum is an aluminum salt (Aluminum potassium sulfate, AIK(SO4)2) and has traditionally been the most widely used adjuvant in commercial vaccines. It has been used for nearly a century as an effective adjuvant. In direct comparisons, higher doses of alum are superior to most other adjuvants, including in

²³ <u>https://jamanetwork.com/journals/jama/fullarticle/2793517</u>

²⁴ <u>https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0035421</u>

²⁵ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4337527/</u>

intranasally delivered vaccines. Alum has been found to be equal or superior to most others for eliciting systemic immunity (in IgG1, IgG2a, IgG2b), and superior for eliciting secretory IgA (sIgA) mucosal immunity and protection against virus challenge in animal studies.²⁶ Moreover, seroconversion in response to vaccination becomes more difficult with age; thus, older people might benefit from the addition of additional potent adjuvants, including those that promote some degree of Th2 response. This is an important consideration for SARS-CoV-2, since older people are most susceptible.

Many other adjuvants can be considered. These include chitosan, inactive cholera toxin (CT), CpG DNA, monophosphoryl lipid A (MPL), poly (I:C), imiquimod and related compounds, and *E. coli* Heat Labile Toxin. Nonetheless, alum has been found to be equal or superior to most others for eliciting systemic immunity (in IgG1, IgG2a, IgG2b), and superior for eliciting secretory IgA (sIgA) mucosal immunity and protection against virus challenge in animal studies.²⁷ Poly (I:C) can act as an additional adjuvant; however, it shows weak activity relative to alum, and at higher doses it has triggered severe reactions and autoimmunity in animal models.²⁸

Chitosan-based therapeutics have been developed for many biomedical applications. Chitosan is a partly or completely deacetylated derivative of chitin, the linear polysaccharide found in the shells of crustaceans like shrimp, exoskeletons of insects, and in fungi. Chitosan nanoparticles were first developed in the mid 1990s by groups in Japan and Spain for drug delivery²⁹,³⁰. For vaccine applications, chitosan acts as both delivery nanoparticle and adjuvant and has been formulated for intranasal administration, parenteral injection, oral and sublingual delivery, and more. It has been used widely in animal experiments, and it has been used safely in human clinical and preclinical trials (see Smith et al. review of trials by El-Kamary et al. and Atmar et al.³¹) (some trials used the commercial product Viscogel)³²,³³ (also see selected human trials reviewed by Guro Gafvelin and Hans Grönlund, in *Molecular Vaccines: From Prophylaxis to Therapy -Volume 2*, Edited by Matthias Giese; Chapter 39, pp. 624-629; Springer. See Table 39.1). The use of chitosan in intranasal delivery of an influenza vaccine to healthy human volunteers produces systemic (IgG) protection, though less than parenteral vaccine with alum adjuvant (although 15 microgram dosage is low)³⁴; plus, it also induces mucosal immune response, unlike parenteral

²⁶ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3868302/</u>

²⁷ https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3868302/

²⁸ <u>https://www.ncbi.nlm.nih.gov/pubmed/?term=32122615</u>

²⁹ <u>https://pubmed.ncbi.nlm.nih.gov/8383199/</u>

³⁰ <u>https://onlinelibrary.wiley.com/doi/10.1002/(SICI)1097-4628(19970103)63:1%3C125::AID-APP13%3E3.0.CO;2-4</u>

³¹ https://www.tandfonline.com/doi/full/10.4161/hv.27449

³² <u>https://cordis.europa.eu/project/id/261954/reporting</u>

³³ <u>https://pubmed.ncbi.nlm.nih.gov/25218298/</u>

³⁴ <u>https://pubmed.ncbi.nlm.nih.gov/15916838/</u>

vaccine³⁵. Chitosan is a cationic polymer typically insoluble at physiological pH, which has led to the development of a range of more soluble derivatives, including trimethyl chitosan (TMC) and quaternary or quaternized chitosan, including N-(2-hydroxy)propyl-3-trimethyl ammonium chitosan chloride (a.k.a. 2-hydroxypropyltrimethyl ammonium chloride chitosans (HTCC or HACC) (for review see Wang et al.³⁶). Cationic chitosan and its derivatives generally are all minimally immunostimulatory and other than physiological pH solubility, their preparation methods are quite similar.

Chitosan as an adjuvant has been compared to alum and to other adjuvants, alone and as part of an adjuvant cocktail, in injectable and intranasal forms. Chitosan alone is a potent mucosal and systemic adjuvant, and it is synergistic with alum and other adjuvants^{37,38}. Importantly, intranasal delivery of chitosan-based vaccines has the ability to elicit both mucosal and systemic immunity.



Schematic showing pathways for Th1 and Th2 acquired immunity (a.k.a. adaptive immunity), and roles of MHC class I and class II. Figure from Akagi, Baba and Akashi³⁹.

³⁵ <u>https://www.sciencedirect.com/science/article/abs/pii/S0169409X01001715</u>

³⁶ <u>https://www.mdpi.com/1422-0067/21/2/487</u>

³⁷ <u>https://pubmed.ncbi.nlm.nih.gov/23933339/</u>

³⁸ <u>https://pubmed.ncbi.nlm.nih.gov/17644205</u>

³⁹ <u>https://link.springer.com/chapter/10.1007/12_2011_150</u>



Th1/Th2 Model for helper T-cells. An antigen is ingested and processed by an Antigen Presenting Cell (APC). It presents fragments from the antigen to T-cells. The upper, Th0, is a T helper cell. The fragment is presented to it by MHC2. IFN- γ , interferon γ ; TGF- β , transforming growth factor β ; mø, macrophage; IL-2, interleukin 2; IL-4, interleukin 4. From Wikipedia⁴⁰.

Chitosan alone elicits both Th1 and Th2 responses, whereas alum triggers strong Th2 and very weak Th1 responses (see Moran et al. for review⁴¹). In general, the Th1 response plays a dominant role in adaptive immunity (a.k.a. acquired immunity) to viral infections, and is responsible for cellular immunity and killing of infected cells by cytotoxic T lymphocytes (CTL), whereas Th2 is humoral and promotes antibody production. Ideally, both Th1 and Th2 processes are elicited by appropriate antigens and adjuvants, and which to target by vaccination are likely to be pathogen specific. Spike protein stimulation of peripheral blood mononuclear cells (PBMC) of SARS-CoV-2 patients results in secretion of high levels of primarily Th1 cytokines, and elevated but much lower levels of Th2 cytokines ⁴². This suggests that SARS-CoV-2 infection provokes a primarily Th1 T-cell response, and a protective vaccine should similarly produce a primarily Th1-based prophylactic response.

⁴⁰ <u>https://en.wikipedia.org/wiki/T_helper_cell</u>

⁴¹ <u>https://pubmed.ncbi.nlm.nih.gov/30195140</u>

⁴² <u>https://www.medrxiv.org/content/10.1101/2020.04.11.20062349v1</u>



Model of chitosan nanoparticle (CSP)-induced NLRP3 inflammasome activation and cross-presentation: Ag together with CSP (via adsorption or encapsulation) was internalized by APCs. The protonation of the amino groups ("proton sponge effect") leads to an extensive inflow of ions and water into the lysosome, which causes the osmotic swelling and deconstruction of the lysosome. The entrapped components (CSP and Ag) are released and finally presented onto MHC I, by cytoplasm degradation (with proteasome and ER involved); After the rupture, lysosome enzymes, cathepsin B, was also leaked into the cytoplasm, which induced the assembly and activation of NLRP3 complex. The capacity of TLR4 stimulation of CSP also played an important role in the intracellular synthesis of pro IL-1β and triggered the secretion of IL-1β and inflammatory responses, together with NLRP3 activation. (Ag: antigen; CSP: chitosan particle.) Figure and caption from Xia et al.⁴³

The amount of vaccine delivered is important. Too little antigen is unlikely to elicit a strong immune response, but too much might attenuate response. Delivery route is also important, with oral delivery being the route most associated with tolerance. Furthermore, high doses of mucosal antigens appear to be required in humans (50+mg) for inducing systemic tolerance, whereas low doses do not appear to induce systemic tolerance.⁴⁴

⁴³ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7185844/</u>

⁴⁴ https://academic.oup.com/intimm/article/26/9/517/2950826

One challenge for mucosal vaccines is verifying systemic efficacy. Many published reports show that nasally delivered vaccines have high efficacy for prevention of infection at the nasal mucosa, typically equivalent or superior to injected vaccines for this most important measure. However, efficacy is not as easily measured or predicted by traditional measures, such as anti-virus or anti-epitope antibody titer (e.g. as measured by ELISA) in serum. This is in part because blood is more quantifiable than mucosal secretions, and in part because the primary means of conferring immunity is through mucosal stimulation and response, which less reliably translates to systemic immunity. Measuring efficacy against SARS-CoV-2 is also difficult because the B-cell/antibody response to coronaviruses is highly variable between individuals, and provides uncertain immune protection. Sustained immune protection appears to be mediated in large part by the T-cell response, which is much more difficult to measure than the antibody response. In certain cases, nasal vaccination can impart mucosal immunity, but no measurable systemic immunity. In such cases, resistance to viral infection due to mucosal immunity will not be predicted accurately by a negative result for IgG systemic antibody. Antibody can be measured in nasal wash, however this is less quantitative and reliable than measurement in serum. Because of such issues, alternative approaches to predicting overall immunity have been developed and published. These include systems biology approaches of blood-based profiling of immune proteins and transcriptomics.

Summary of key general features of an open-source, citizen-science vaccine:

- Safe ingredients. Long history of published results.
- Ease and robustness of production. Simple to make but with the possible tradeoff of short stability. Nanoparticles of chitosan or its derivatives form spontaneously and reproducibly, over a wide range of conditions and ingredient concentrations.
- Needle-free delivery. Intranasal administration is a preferred delivery mode for vaccines against respiratory and nasal viruses, and offers one of the highest safety profiles. Potentially stimulates both protective mucosal immunity at the site of infection, and systemic immunity.
- **Peptide epitope antigens.** Short peptides of linear epitopes are easiest to procure or produce, and are predicted to function best without specific structural constraints. These can be mined from a rich literature, or produced de novo. Antigens for incorporation into a vaccine can be produced synthetically or as recombinant expression proteins.
- **Booster schedule.** Allows use of intranasal delivery and a lower dose or milder adjuvant, yet has the potential to produce immune responses equivalent to a single dose of hyper-stimulatory adjuvanted adjuvants.

POSSIBLE MECHANISMS OF VACCINE-ENHANCED DISEASE, VACCINE-INDUCED AUTOIMMUNITY, AND MITIGATION STRATEGIES (updated 2020-10-24)

As mentioned briefly in the section above, a small number of injected vaccines have led to vaccine-enhanced disease (VED), meaning that disease is made more serious in people or experimental animals that have been vaccinated, relative to unvaccinated controls. This has occurred in response to vaccines for respiratory syncytial virus (RSV), dengue, Zika, and others. Some of these cases involve experimental vaccines, although RSV and dengue were deployed in healthy human populations. Although there isn't a single accepted or proven mechanism known to cause or contribute to VED, there are historical vaccine design features and adjuvant factors that are suspected contributors. The RSV vaccine that caused VED in the 1960s was formalin inactivated whole virus, adjuvanted with alum. This vaccine triggered a weak initial immune response, and theoretical explanations for how it caused VED have focused on the chemical alteration of antigenic structures caused by formalin and on the Th2 bias induced by alum. Recent studies have shown that, in addition to Th2 bias, RSV VED is driven by a few factors including deficiencies in Toll-like receptor activation and CD8 response, and hyperactivation of a distinct subset of CD4 T-cells.^{45,46,47} Mitigation of any of these reduces or obviates VED. With dengue virus, a second infection with a distinct dengue serotype often causes more severe disease than the first infection. Use of the dengue vaccine Dengvaxia similarly results in more severe disease than in unvaccinated controls, suggesting that vaccination sets up a serious subsequent infection by similar mechanisms to a prior dengue infection. (It is important to note that although named "vaccine" enhanced disease, the same phenomenon of enhanced disease with subsequent infection has been described in natural infection with multiple pathogens, and could be due to any number of immune-evasive strategies viruses have evolved.)

One proposed mechanism for VED in dengue is antibody-dependent enhancement (ADE), in which antibodies of the systemic immune system act in "Trojan horse" fashion to facilitate viral entry into certain host immune cells. (For a review of ADE specific to SARS-CoV-2 see Eroshenko et al.⁴⁸) Waning or weak antibody response, including a substantial non-neutralizing antibody component to this response, has been shown to facilitate viral entry into macrophages and monocytes. There are multiple possible ways this might occur. There are multiple dengue serotypes, which are distinguished by substantial protein sequence divergence. Amino acid differences in key epitopes can alter binding of antibodies, and the result is that antibodies that

⁴⁵ <u>https://www.nature.com/articles/nm.1894</u>

⁴⁶ <u>https://pubmed.ncbi.nlm.nih.gov/25769044/</u>

⁴⁷ <u>https://pubmed.ncbi.nlm.nih.gov/17911628/</u>

⁴⁸ <u>https://www.nature.com/articles/s41587-020-0577-1</u>

are neutralizing for one serotype bind antigens of another serotype, but in a manner that is weakly or non-neutralizing. One possible contributing factor in this process is a phenomenon referred to as "original antigenic sin," or "antigenic imprinting." Exposure to an initial antigen by infection by one serotype produces memory immune cells that recognize the amino acid sequence of this antigen, and upon subsequent exposure to a related but slightly different antigen, there is preferential expansion of memory cells that recognize the original antigen rather than cells that recognize and bind more tightly to the second antigen.⁴⁹ Thus, upon second infection, the antibody response is directed mostly at epitopes of the previous infection, and less robustly against the current infection. Antigenic imprinting is not limited to B-cell epitopes, and essentially the same phenomenon has been reported for T-cell epitopes and responses.⁵⁰

Other aspects of the T-cell response also have been implicated in VED. In particular, a robust CD8 response is a key immune correlate of mild or asymptomatic infection for a number of respiratory viruses, including dengue, influenza, SARS-CoV-1, and SARS-CoV-2. Multivalent whole-virus dengue vaccines that provide robust and broad protection against dengue infection produce a robust CD8 response.⁵¹ Notably, the CD8 epitopes that produce the strongest protection against all dengue serotypes are derived from conserved non-structural proteins, rather than the viral outer surface proteins, which have much greater amino acid sequence variation.⁵² This appears to be a common feature of respiratory viruses, including influenza and SARS-CoV-2.^{53,54}

In contrast, Dengvaxia is based on a synthetic hybrid virus that uses the non-structural backbone of the related yellow fever virus, together with the outer structural proteins from dengue. Dengvaxia elicits a strong initial antibody response, but this strong response wanes relatively quickly. Since Dengvaxia lacks the immunodominant CD8 epitopes of the non-structural proteins, the cytotoxic response is directed toward the weakly immunogenic dengue surface epitopes, and the immunodominant non-structural proteins of the yellow fever virus. Therefore, not only is the CD8 response specific to dengue epitopes relatively weak, but a possible exacerbating factor is that the primary CD8 response triggered by Dengvaxia is directed away from dengue by immunodominant yellow fever epitopes. To summarize one model of Dengvaxia-induced VED favored by some leading immunologists: upon waning antibody response (and possible ADE), compromised cellular immunity is weak, and upon subsequent infection T-cells mount a pathological cellular immune response, possibly exacerbated by antigenic imprinting, driven by

⁴⁹ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7291596/</u>

⁵⁰ <u>https://pubmed.ncbi.nlm.nih.gov/9697771/</u>

⁵¹ <u>https://jvi.asm.org/content/89/1/120.full</u>

⁵² https://www.frontiersin.org/articles/10.3389/fimmu.2019.02125/full

⁵³ https://pubmed.ncbi.nlm.nih.gov/18802496/

⁵⁴ <u>https://www.nature.com/articles/s41590-020-0782-6</u>

the highly variable surface T-cell epitopes of the dengue serotype of the post-vaccination infection.

It is important to reiterate that immunopathological responses are not limited to vaccines, and are known to be mediated by viral exposures, as occurs in primary dengue virus infection followed by second infection by a distinct dengue serotype. However, as seen in RSV and dengue, vaccine designs can contribute to the likelihood of immunopathology. One early report on SARS-CoV-2 suggests that neutralizing IgG antibodies for the SARS-CoV-2 receptor binding domain (RBD) do not exhibit such enhancement; and as mentioned above, some experts suggest that ADE is not a clear concern in the development of vaccines against this virus^{55,56}. However, other experts are less confident that SARS-CoV-2 vaccine-induced ADE or other VED will not occur.

Given the immunopathology associated with Dengvaxia, we are especially concerned about the use of the same general design features that are suspected to have caused or contributed to VED. In particular, if SARS-CoV-2 immunodominant MHC class I/CD8 epitopes are concentrated on non-surface proteins, use of SARS-CoV-2 outer surface proteins alone might produce a weak CD8 response; and if this feature is combined with the backbone of another virus carrying potentially immunodominant CD8 epitopes, such a vaccine fits a general risk profile similar to Dengvaxia. As of late 2021, the most extensive quantitative T-cell epitope mapping data were published by Adaptive Biotechnologies and made public in the ImmuneCODE database⁵⁷. The mapping of SARS-CoV-2 epitopes shows clear CD4 immunodominance, and also some degree of CD8 immunodominance. It is notable and concerning that the majority of immunodominant MHC class I/CD8 epitopes (7 of the top 8; 12 of the top 15) of the entire proteome are found in non-structural proteins, analogous to the distribution found in dengue.

Vaccine-induced autoimmunity (VIA) is another immunopathological complication of vaccination. It is a rare but worrisome side effect. Over the past decades, several vaccines have been implicated as autoimmune triggers⁵⁸. Four possible factors have been proposed in explanations for mechanisms of VIA: molecular mimicry and cross-reactivity to host antigens; unbalanced or hyperstimulation of immunity by specific adjuvants; host genetics; and concurrent vaccination and infection. A combination of molecular mimicry of host antigens and certain potent adjuvants is especially concerning, and has been suggested as a leading explanation for VIA in narcolepsy triggered by Pandemrix, although specific Scandinavian host genetics appear to play a role.

⁵⁵ <u>https://www.biorxiv.org/content/10.1101/2020.04.10.036418v1</u>

⁵⁶ https://www.nature.com/articles/s41587-020-0577-1

⁵⁷ https://www.researchsquare.com/article/rs-51964/v1

⁵⁸ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6078966/</u>

Other serious autoimmune diseases believed to be triggered by vaccines are Guillain-Barré syndrome, Multiple Sclerosis, and lupus. While these side effects are quite serious, they are also quite rare. The incidence of narcolepsy increased over 10-fold in those vaccinated with Pandemrix, but the vaccine-attributable rate was only 1 in 18,400⁵⁹. In each case, molecular mimicry is a suspected primary contributor, and viral infection can trigger these same autoimmune reactions, sometimes even more frequently than vaccines. Therefore, as with other decisions regarding vaccination during a global pandemic, if certain antigens pose an increased risk of autoimmunity, one is likely to be exposed to those antigens eventually either by viral infection, or through vaccination. And in the case of vaccination, the overall adverse-outcome risk profile may be far lower.

In the list below, we summarize possible mechanisms of VED in precedent cases that might contribute to VED immunopathology in response to vaccination against SARS-CoV-2, along with some of the mitigation strategies we have used to address each of these in our vaccine designs.

- Th2 and/or Th17 immune hyperpolarization, possibly accompanied by an attenuated Th1 response have been proposed to be leading risk factors for VED. It has been suggested that use of alum as an adjuvant creates such a Th2 bias and inhibition of Th1, while the use of other adjuvants, such as chitosan and toll-like receptor (TLR) agonists shift the balance in favor of Th1.
 - a. In many published studies of chitosan, our chosen delivery and adjuvant vehicle, intranasal delivery elicits a balanced Th1/Th2 immune response.
 - b. TLR agonists have been used together with chitosan, including in intranasal formulations to further enhance the Th1 response. In order to simplify vaccine production and to create a higher margin of short-term safety, we have not included additional adjuvants in our designs; however, the inclusion of such secondary adjuvants is relatively straightforward, and might be used to further increase the Th1:Th2 ratio.
- ADE of viral infection is primarily associated with IgG and the Fcγ receptor, and much less often with IgA and IgM, and their Fc receptors.⁶⁰ Secretory IgA (sIgA) of mucosa has been shown to mediate ADE of bacterial infection but has not been demonstrated for viral infection.⁶¹
 - a. Secretory IgA is the dominant antibody isotype at mucosal surfaces. It remains unknown if ADE of viral infection can occur through sIgA, but it has not been reported. Nevertheless, a robust mucosal immune response should greatly reduce

⁵⁹ https://pubmed.ncbi.nlm.nih.gov/28847694/

⁶⁰ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7119964/</u>

⁶¹ <u>https://pubmed.ncbi.nlm.nih.gov/12967664/</u>

or prevent IgG-mediated ADE in the systemic response by reducing or abrogating initial infection.

- Certain B-cell epitopes appear to be more likely to give rise to ADE, including on the Spike protein of SARS-CoV-1, resulting in concern about the use of full-length Spike protein in SARS-CoV-2 vaccines.⁶²
 - a. Spike is the primary target for generating antibody-based protective immunity. It is a very large protein (1273 amino acids) with many antibody binding epitope sites, but only a small number of prospective epitopes are potentially neutralizing. This has been proposed to be a potential contributing factor to ADE.
 - b. Selected target epitopes included in the vaccine represent only a fraction of epitopes, including those shown to be bound by neutralizing antibodies. In the current vaccine design, only about 5% of the Spike protein sequence is represented as B-cell epitopes. If original antigenic sin occurs, it should only apply to these epitopes, and not affect other neutralizing epitopes of the virus or subsequent vaccines. This allows epitopes of this subsequent exposure to trigger activation, maturation and expansion of naive B-cells into plasma cells that secrete antibodies specific to these neutralizing epitopes of the second infecting viral serotype. There are multiple Spike protein conformational B-cell epitopes bound by neutralizing antibodies, and peptide vaccines such as those described in this white paper series do not include many of these portions of Spike.
 - c. Incorporation into a vaccine of only a small portion of all viral antigens also reduces the likelihood that a viral antigen will trigger VIA through molecular mimicry and immune cross-reactivity to a host antigen. Again, during a pandemic, if certain antigens of a pathogen pose an increased risk of autoimmunity, one is likely to either be exposed to antigens by viral infection, or through vaccination. Selected epitope peptide sequences can be compared to the human proteome (e.g. by BLAST) to further reduce cross-reactivity to host antigens. We perform a BLAST for every selected epitope peptide to minimize the potential for high-scoring matches to potentially cross-reactive host antigens.
- 4. Waning or weak T-cell response, possibly driven by T-cell cross-reactivity to divergent epitopes derived from prior exposures to different viral serotypes. One model for how this might occur is through a T-cell original antigenic sin, analogous to the phenomenon described for antibodies.⁶³
 - a. Again, at the scale of a large population, it is impossible to completely avoid this risk since it is not restricted to vaccines, but also applies to viral infection.

⁶² https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2756483/

⁶³ https://www.jimmunol.org/content/176/6/3821.long

Nevertheless, selection of a few conserved epitopes should reduce the likelihood that a SARS-CoV-2 mutant will arise that results in this phenomenon. Following the same logic described for B-cell epitopes, if original antigenic sin occurs it can be limited to only those epitopes supplied in the vaccine. In the current vaccine design, less than 1% of the SARS-CoV-2 proteome (entire protein sequence) is represented as T-cell epitopes, and only about one-quarter to one-third of empirically verified immunodominant epitopes.

- 5. Similar to SARS-CoV-2, severe dengue infection is CD8 deficient, while mild infection is accompanied by a robust CD8 response. Multivalent dengue vaccines that elicit protective immunity include immunodominant MHC class I/CD8 epitopes from non-structural proteins. The Dengvaxia design employs a hybrid virus that uses the non-structural backbone of the related yellow fever virus, together with the outer structural proteins from dengue. Immunodominant MHC class I/CD8 epitopes are concentrated in the non-structural proteins of dengue, SARS-CoV-2, and other respiratory viruses, while immunodominant MHC class II/CD4 epitopes are concentrated in the outer structural proteins. Dengvaxia elicits a strong initial, but rapidly waning antibody response, and a weaker and possibly immunopathological CD8 T-cell response in response to subsequent infection.
 - a. RaDVaC vaccine designs incorporate a few immunodominant MHC class I/CD8 epitopes from non-surface and non-structural proteins. Some of these epitopes are highly conserved, and the most common HLA alleles are covered redundantly by at least 2 separate epitopes. We have not included all immunodominant epitopes for CD8 (or for B-cells or HLA class II/CD4), which should prevent original antigenic sin from compromising a future robust immune response to a cross-reactive serotype.
- 6. VED might arise through vaccine design-induced distortion of viral epitopes, as has been proposed to occur by formalin treatment of RSV.
 - a. Our preferred epitope type is synthetic peptides. Such peptides are chemically well defined and have a high degree of structural integrity.

KEY TECHNICAL FEATURES AND SPECIFICATIONS (Updated 2021-03-06)

Certain technical features of the RaDVaC strategy follow from the above considerations, and are key to an open-source, citizen-science vaccine.

1. **Delivery.** Intranasal or sublingual delivery are probably among the safest choices of administration routes, though other delivery modes, especially other mucosal delivery

modes, are likely also safe. For a respiratory virus, both intranasal and sublingual delivery have the advantage of eliciting a mucosal immune response at the most common sites of viral entry. These delivery modes can also elicit systemic immunity, although often not as robust as parenteral delivery after a single dose. Mucosal delivery is also less likely than other routes to trigger vaccine hesitancy.

2. Chitosan nanoparticle and adjuvant(s). Successful vaccines depend on successful delivery and on immunogenic stimulation, often by the use of adjuvants that promote appropriate stimulation of the immune system. Many adjuvants are proven and available, with various strengths and weaknesses. A leading nanoparticle combined nasal delivery and adjuvant system is based on chitosan and sodium triphosphate (STP) a.k.a. tripolyphosphate (TPP)⁶⁴, which is the present top choice for this intranasal vaccine. Chitosan nanoparticles of various sizes can be created with simple tunable parameters.⁶⁵,⁶⁶ The virus has an average size of about 100 nm to 150 nm.⁶⁷ Nanoparticle size ideally will be tuned to be between 100 nm and 200 nm. As described above, chitosan has been shown to be safe and well tolerated, and intranasal delivery elicits both mucosal and systemic immune responses. Chitosan is available in various molecular weights, deacetylation ranges, and forms such as pure chitosan, trimethyl chitosan, and quaternary or quaternized chitosan, including hydroxypropyl-trimethyl ammonium chitosan chloride (HTCC or HACC). It is important to obtain high purity chitosan (including a certificate of analysis). Low-grade chitosan might not work well, or might be contaminated with allergenic shellfish proteins. Multiple forms, molecular weights and suppliers have been tested and published. Very low molecular weight chitosans (e.g. 1 to 20 kilodaltons; sometimes referred to as oligochitosan) have the advantage of being water soluble, yet the lower the molecular weight, the less stable the chitosan gel. Unmodified higher molecular weight (100+ kDa) chitosans are only soluble at pH values under their pKa values of 6.2 to 6.5 because their free amino groups are protonated. Deprotonation at physiological pH causes insolubility and loss of integrity of nanoparticles formed by ionic crosslinking. Trimethyl chitosan and HACC do not suffer from this problem, because they possess amino groups that are protonated at physiological pH. Therefore, after much experimentation we currently use HACC as our primary chitosan derivative, although some RaDVaC collaborators (aka Radvackers) are experimenting with trimethyl chitosan. Throughout this document, the general term chitosan can mean unmodified chitosan along with soluble derivatives, such as HACC and trimethyl chitosan.

⁶⁴ https://www.frontiersin.org/articles/10.3389/fimmu.2018.00934/full

⁶⁵ https://www.nature.com/articles/s41598-018-23064-4

⁶⁶ <u>https://www.ncbi.nlm.nih.gov/pubmed/28578269</u>

⁶⁷ https://www.nejm.org/doi/full/10.1056/NEJMoa2001017

- a. Possible additional adjuvants/immunostimulants. Chitosan is a Th1/Th2 immune triggering, self-adjuvanting polysaccharide; therefore, additional adjuvants or immunostimulants are not strictly necessary, and none will be incorporated into the initial formulation. However, if chitosan alone produces insufficient immune stimulation, we are considering experimenting with the incorporation of other adjuvants that enhance a Th1 or specific T-cell targeted response. Good candidates are alum (Th2 biased), imiquimod (TLR7 agonist, Th1 biased), all-trans retinoic acid (ATRA)⁶⁸, poly (I:C) derivatives (TLR3 agonist, Th1 biased), compound 48/80 (mast cell agonist with balanced Th1/Th2/Th17 response)⁶⁹, and other TLR⁷⁰ (mostly Th1 biased) and mast cell⁷¹ agonists.
- 3. **Peptide antigen.** A peptide-based vaccine is the method of choice, based on and bearing in mind the following considerations:
 - Multimeric. Multimeric/multi-epitope vaccines have been shown to be effective. The approach taken here uses multiple peptides, each of which will carry one or more B and/or T-cell epitopes.
 - b. **Fast and flexible**. Synthetic peptides are inexpensive, and can be made to order quickly by many commercial peptide producers. They can be synthesized with many possible chemical modifications that have been reported to increase immunogenicity.
 - c. Epitope conformation and presentation. Gen 10+ designs feature surface display of B-cell epitope peptides for enhanced antibody production. Early versions of the RaDVaC vaccine contained simple linear epitopes, with no special regard to conformation or presentation. Some of the B-cell epitope peptides from Gen 3 onward were conformationally constrained with disulfide bonds. Ideally, 3D structures of viral proteins should be imaged and linear epitopes should be chosen that either don't require special conformational constraints, or attempts should be made to achieve natural structural conformation of the epitope peptide.
 - d. **Non-peptide antigens**. Other antigens are possible and can be delivered by chitosan intranasally, including proteins, DNA or RNA.
- 4. **Epitope selection** is critical. B-cell and T-cell epitopes have been selected and published by others. Multiple epitopes of both types should be selected, preferably in important functional and/or conserved regions of the virus. This gives a higher probability that at

⁶⁸ https://jvi.asm.org/content/85/16/8316

⁶⁹ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2743390/</u>

⁷⁰ https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3000864/

⁷¹ <u>https://www.nature.com/articles/nm1757</u>

least one epitope will successfully stimulate immunity, especially given variability in individual immune responses, rather than relying on a single epitope.

- a. B-cell epitopes. Studies of antibody mapping from convalescent patients help identify portions of the virus available for antibody binding to B-cell epitopes, and some of these antibodies have been shown to neutralize viral infection in cellular studies or specially designed neutralization assays. Certain B-cell epitopes that score highly by common machine learning predictions have not been reported at high frequency in convalescent sera. This might be due to linear peptide-based approaches used to map epitopes, the extensive glycan shield around the virus, or other complicating factors. Whatever the explanation, we regard the convalescent antibody data as superior to purely computational predictions.
- b. T-cell epitopes. Empirical methods similar to but more complex than those used for B-cell epitope selection have been used for selecting superior T-cell epitopes.
- 5. Testing. Success of mucosal and systemic humoral immunogenic stimulation will be assessed by testing antibody titers of nasal wash, saliva, and serum. Tests of cellular immunity (T-cell) will also be performed, although as of mid Feb 2021 there are no commercial options yet available for these kinds of tests, which require substantial laboratory infrastructure. Testing of immunity should be performed using standard assays and reagents, as well as the use of newer technologies, such as transcriptome profiling of peripheral blood mononuclear cells (PBMCs).

GOALS (last updated 2022-01-14)

- 1. Design and incrementally improve a vaccine made with commercially available materials, and with the following properties:
 - All ingredients have undergone extensive testing for efficacy and safety in animal models, and preferably in human trials. Delivery and adjuvant materials and technologies must have decades of proven safety.
 - b. Use of best practices to achieve high probability of efficacy.
 - c. Mode of administration must be safe and proven. Intranasal delivery is preferred for safety and for establishment of both mucosal and systemic immunity. Inhalation and injection are possible delivery modes for this vaccine but should only be pursued by those with sufficient training and experience.
 - d. Vaccine should be extremely easy to make (including sourcing of ingredients) and administer.
- 2. Produce vaccine for initial testing, and self-administration.

- 3. Produce and administer updated generations of vaccines as new information becomes available.
- 4. Design and implement testing for immune responses to the vaccine. Tests will be performed with published methods, and new methods should be developed to predict efficacy. Ideally, biological samples such as blood and nasal wash are collected before initial vaccination, and at various time points.
- 5. Openly share and disseminate information, enabling many others to produce the vaccine. The only custom components are the epitope peptides. By themselves, these synthetic peptides are essential components not only of vaccine, but also of standard types of testing (ELISA, ELISpot, etc.), and thus are not drugs, and might be distributed to facilitate progress in multiple kinds of research.
- 6. Maintain ongoing long-term self-monitoring of health and symptoms to collect empirical evidence of degree of efficacy and possible unforeseen effects.
- 7. Create an online forum for the exchange of data, ideas and best practices with other individuals and groups. Over time, results can be aggregated for broader analysis.
- 8. Create and openly share tools for design and production of vaccine components. For example, openly licensed recombinant organisms capable of producing viral subunits usable for vaccines.

MATERIALS AND METHODS (updated 2021-03-07)

Selection and purchase of the vaccine ingredients

All materials and ingredients are commercially available. However, some vendor deliveries might be interrupted by public health measures. The vaccine is very simple and consists of five ingredients: epitope peptide(s), chitosan, sodium triphosphate, water, and sodium chloride.

- Epitope peptides. Multiple vendor options are available for synthetic peptides. Peptides or concatemers of the selected epitopes can also be produced by other means, such as recombinant plasmid-based expression and purification (e.g. E. coli His tag). Options should be considered and tested, as supply channels might be disrupted by the pandemic.
- Chitosan derivatives. Chitosan is available in various molecular weights, deacetylation ranges, and forms such as pure chitosan, trimethyl chitosan, and quaternary or quaternized chitosan, including N-(2-hydroxy)propyl-3-trimethyl ammonium chitosan chloride (HTCC or HACC, also known as N-HTCC or N-HACC). (Note that O-HTCC or O-HACC are not the same as N-HACC, and degrees of substitution and therefore

<u>molecular weights will differ, although they likely will work similarly.</u>⁷² It is important to obtain high purity chitosan (including a certificate of analysis). Low-grade chitosan might not work well, or might be contaminated with allergenic shellfish proteins. Because unmodified chitosan is water insoluble at physiological pH, undermining nanoparticle integrity *in vivo*, we currently use HACC as our primary chitosan derivative (although some Radvackers are experimenting with trimethyl chitosan). Acceptable ranges: medium to high molecular weight range of 100-400 kDa. Deacetylation = 75% to 90%. Depending on the source, some initial experimentation is likely necessary to test various behaviors of the stock solution. If you only have access to unmodified chitosan, pre-Gen 10 protocols can be used⁷³; however, soluble chitosan derivatives will give superior results.

- Sodium triphosphate, Na₅P₃O₁₀. Multiple suppliers.
- Sodium chloride (NaCl). 5M stock solution.
- dH₂O, deionized water. Widely available.
- OPTIONAL (If you are using unmodified chitosan instead of soluble chitosan derivative): Distilled white vinegar (or acetic acid CH₃COOH). White vinegar is typically about 5% acetic acid.
- OPTIONAL: Granular activated charcoal for circularizing optional peptides.
- OPTIONAL: Sodium bicarbonate NaHCO₃, or other mild base for adjusting the pH.

Tools for designing and analyzing vaccine components (NEW 2022-07-22)

Here are a few useful tools for analyzing epitope peptides and sequence data. There are many more we use, and we'd like to hear of tools you like in order to share useful information with the RaDVaC network.

- Immune Epitope Database and Analysis Resource (IEDB) MHC-I Binding Predictions, MHC-I Binding (iedb.org)
- NetMHCpan server for analyzing MHC/HLA types, <u>https://services.healthtech.dtu.dk/service.php?NetMHCpan-4.1</u>
- A Plasmid Editor (APE), <u>https://jorgensen.biology.utah.edu/wayned/ape/</u>
- Peptide synthesis analysis tool (Thermo Fisher),
 <u>https://www.thermofisher.com/us/en/home/life-science/protein-biology/peptides-proteins/</u>
 <u>custom-peptide-synthesis-services/peptide-analyzing-tool.html</u>
- Peptide solubility calculator (Innovagen),
 <u>https://pepcalc.com/peptide-solubility-calculator.php</u>

⁷² <u>https://onlinelibrary.wiley.com/doi/pdf/10.1002/app.26038</u>

⁷³ <u>https://radvac.org/wp-content/uploads/2020/12/White-Paper-SARS-CoV-2-vaccine-ver-3-3-1.pdf</u>

Preferred equipment and materials (updated 2020-11-04)

Certain equipment is required for efficient vaccine production and administration. Vaccine can be produced without specialized laboratory equipment but the process is more laborious and the results are likely to be more variable.

- Pipettes: 1000 microliter, 200 microliter
- Sterile filtered pipette tips: 1000 microliter, 200 microliter
- Scale: gram scale accurate to 0.1 grams, or jeweler's scale for very small amounts
- Clean spatula for dry reagents
- Small beakers for mixing and stirring. 10 ml to 25 ml beaker for stirring
- Small bottles for stock solutions; 100 ml to 250 ml
- 15 ml conical tubes
- 50 ml conical tubes
- 1.5 to 2 ml microcentrifuge tubes
- Stirring apparatus: magnetic stir plate and small stir bar to fit in beaker
- OPTIONAL but preferable: pH strips, range at least 2.0 to 9.0
- Nasal spray apparatus. These are available through multiple vendors. We selected small
 nasal spray bottles (about 5 milliliter) with a pump-top apparatus. A tube stem about 4 cm
 long and approximately 1 mm inner diameter extends from the bottom of the apparatus,
 which can be placed into the bottom of a 2-ml tube containing vaccine. Our spray
 apparatus delivers about 100 microliters per pump.

Selection of epitopes (updated 2021-03-17)

Epitopes are a critical element of a vaccine because they are specific to each pathogen. An epitope is the portion of an antigen (typically a protein) that interacts with the immune system, such as through binding of antibodies or T-cell receptors. Multiple data types can be used in the selection of epitopes. This section describes general guidelines and issues for consideration in epitope selection and design.

- Epitopes for B-cell and T-cell responses can be quite different, and there are critical differentiating features. Here are key considerations in the selection and design of B-cell and T-cell epitopes.
 - T-cell epitopes are presented as short linear peptide antigens, but B-cell epitopes (which can be linear or conformational) are presented in their native conformation.
 - The B-cell response is relatively easily measured by antibodies, but the T-cell response is more difficult to measure and more rarely performed. Antibody testing

is a large-scale commercial activity (using lateral flow, ELISA, etc.) but T-cell testing is generally a research activity (using ELISpot, etc.). Therefore, T-cell epitopes are often based on computational prediction. This was true of early T-cell epitope selections in the biomedical literature and preprint servers. However, beginning in mid July, 2020, substantial empirical data have been published on T-cell epitopes.

- B-cell or antibody binding epitopes are often mapped into two general classes: linear and conformational (or structural). Linear epitopes (single peptide chain of nearby amino acids) are typically mapped by binding antibody-containing serum from convalescent patients to large collections or arrays of short epitope peptides collectively representing viral proteins. Conformational epitopes often consist of amino acids that are far apart on a protein sequence but in close proximity in the folded protein. Linear epitopes are easier to map than conformational epitopes. However, conformational epitopes can be critically important.
- There are 2 classes of T-cell epitopes: MHC Class I and MHC Class II (for a thorough review, see Rock et al.⁷⁴). Core motif binding of both Class I and Class II is about 9 amino acids but both are variable. The Class I binding site has closed ends, and epitopes range from 8 to 15 aa, but infrequently diverge from 9 aa. Class II binding is open-ended, and natural epitopes extend longer than the 9 aa core, and range from 13 to 25 aa⁷⁵. Many computational predictive algorithms are available, and a large number of preprints provide Class I and Class II predictions for inclusion in a vaccine. If empirical data are available for protective T-cell epitopes (e.g. Le Bert et al, Adaptive Biotechnologies, Peng et al⁷⁶, and Zhang et al⁷⁷), these data should guide T-cell epitope selection.
- Immune responses (both B and T-cell) can vary substantially with different ancestry/ethnicity. We have used predictions and empirical data for wide coverage of immune responses (our group is highly ethnically diverse). For specific geographies and ancestries, local immune response information should be taken into account.
- Although there can be differences between T and B-cell epitopes, they can overlap. If possible, peptides that contain experimentally validated epitopes for both B and T-cells are ideal.
- Immunodominance and limited effective epitope space. Immunodominance is the phenomenon of antibody and/or T-cell responses to a small number of epitopes, despite

⁷⁴ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5159193/</u>

⁷⁵ <u>https://pubmed.ncbi.nlm.nih.gov/1380674/</u>

⁷⁶ https://www.nature.com/articles/s41590-020-0782-6

⁷⁷ <u>https://pubmed.ncbi.nlm.nih.gov/32668444/</u>

a large number of possible epitopes within the virus. Findings to date suggest both B-cell and T-cell responses to SARS-CoV-2 display pronounced immunodominance. Some key findings about the immunodominant landscape of SARS-CoV-2:

- A publication by Snyder and colleagues shows a few epitope peptides are responsible for over 50% of CD8 T-cell receptor binding events⁷⁸.
- The immunodominance landscapes of CoV-1 and CoV-2 are:
 - Very different between CoV-1 and CoV-2 Spike proteins, possibly largely due to glycosylation differences.⁷⁹
 - Highly similar for CoV-1 and CoV-2 Nucleocapsid proteins^{80,81}.
- One early but thorough preprint by Zhang and colleagues suggests immunodominance is a feature of the B-cell immune response to SARS-CoV-2⁸².
- Farrera and colleagues reported a set of immunodominant B-cell epitopes similar to those reported by Zhang, but in a study of Swiss convalescent patients⁸³.
- Most of the epitopes reported by Zhang et al and Farrera et al were subsequently confirmed by Li and colleagues in the first large-scale linear B-cell epitope mapping of over 1,000 convalescent patients in Wuhan, China⁸⁴.
- The emerging consensus suggests that there are about half a dozen dominant B-cell epitopes, which are the same in European and Chinese patients. The majority are found in key functional portions of the Spike protein, including the following (amino acid coordinates shown are approximate):
 - Spike 450-500; ACE2 binding residues of the RBD (Zhang et al); low degree of conservation; probably moderate to high mutant escape potential
 - Spike 655-685; immediately adjacent to the S1/S2 protease cleavage sites; high degree of conservation in the antibody-binding N terminal portion of the epitope; likely low to moderate mutant escape potential. Cleavage of this site greatly increases efficiency of host cell infection⁸⁵.
 - Spike 805-820; spanning the S2' protease cleavage site (815-816); high degree of conservation; likely low mutant escape potential.

⁷⁸ <u>https://www.medrxiv.org/content/10.1101/2020.07.31.20165647v2</u>

⁷⁹ <u>https://pubs.acs.org/doi/full/10.1021/acsinfecdis.6b00006</u>

⁸⁰ <u>https://jcm.asm.org/content/42/11/5309.full</u>

⁸¹ <u>https://www.biorxiv.org/content/10.1101/2020.03.26.994756v1</u>

⁸² https://www.biorxiv.org/content/10.1101/2020.04.23.056853v2.article-info

⁸³ https://www.medrxiv.org/content/10.1101/2020.06.15.20131391v1

⁸⁴ <u>https://www.medrxiv.org/content/10.1101/2020.07.13.20152587v1.full.pdf</u>

⁸⁵ <u>https://www.pnas.org/content/117/21/11727</u>

- Spike 1145-1160; adjacent to the heptad repeat 2 (HR2) region, which is critical for fusion of the viral and host cell membranes of the SARS virus⁸⁶; very high degree of conservation; likely extremely low mutant escape potential.
- These immunodominance findings are supportive of the peptide vaccine approach, since only a limited portion of the virus needs to be represented in the vaccine for it to be effective; plus, potentially counterproductive portions of the virus (such as those that might contribute to ADE) are not present in the vaccine.
- The critical importance of MHC Class I/CD8 T-cell epitopes and immunity. Antibodies
 get a lot of press attention (in large part because simple commercial antibody tests are
 widely available but T-cell tests are more complex and must be performed on isolated
 cells, and therefore are done in specialized research labs) but the most protective
 immune response consists of robust humoral (B-cell mediated) and cellular (T-cell
 mediated) responses. Early indications suggest T-cell responses might be key to both
 minimizing severity of illness and to longevity of immunity, but the specific type(s) of T-cell
 response that might be most protective is a critical yet still unresolved issue. Here are a
 few key findings about the importance of T-cell immunity to SARS-CoV-2 infection:
 - Mild COVID-19 is seen in convalescents with robust T-cell responses, including those who do not have detectable antibodies (are seronegative)⁸⁷
 - Severe COVID-19 is accompanied by higher antibody levels than mild cases, but with unbalanced or impaired T-cell response, especially the CD8 response⁸⁸. This study by Peng et al identified 6 immunodominant CD4 epitopes (3 of which are located on the Spike protein), a similarly small number of immunodominant CD8 epitopes, and they showed that a higher ratio of CD8/CD4 response is found in those with milder disease versus those with severe disease.
 - The effect of T-cell epitopes can vary depending on the protein of origin. Peng and colleagues also report that T-cell epitopes from Spike, Membrane and ORF3a proteins were associated with more severe disease. Higher frequencies of multi-cytokine producing Membrane and Nucleocapsid-specific CD8 T-cells (relative to Spike-specific CD8 T-cells) were observed in patients with mild cases. Because of these findings, Peng et al conclude that "The identification of T-cell specificity and functionality associated with milder disease, highlights the potential importance of including non-spike proteins within future COVID-19 vaccine design." Many subsequent publications have further strengthened this conclusion.

⁸⁶ <u>https://www.thelancet.com/journals/lancet/article/PIIS0140673604157887/fulltext</u>

⁸⁷ https://www.biorxiv.org/content/10.1101/2020.06.29.174888v1

⁸⁸ <u>https://www.biorxiv.org/content/10.1101/2020.06.05.134551v1.full.pdf</u>

Nevertheless, the CD4 epitopes included in all RaDVaC designs to date do not include the 3 immunodominant CD4 Spike protein epitopes identified by Peng et al, and we have no plan to include these in future designs until this concern about CD4 hyperstimulation is resolved.

- Peng et al also used single cell analysis of bronchoalveolar lavage fluid from COVID-19 patients and found substantial clonal expansions of CD8 T-cells in mild but not severe patients, further reinforcing evidence that a robust CD8 response protects against severe disease.
- T-cell exhaustion or deficiency is correlated with disease severity (low T-cell response with more severe disease) ^{89,90,91,92}
- Population MHC class I binding (and CD8 response) is predictive of mortality⁹³.
- Vaccine-enhanced disease involves a Th2-polarized response that is CD8 T-cell deficient⁹⁴. A robust CD8 response is observed in people and animal models who do not develop vaccine-enhanced disease.
- Many reports have shown rapid declines of anti-SARS-CoV-2 antibodies in convalescents. It remains unclear how this decline impacts protection from reinfection or severity of disease, but initial results also show that T-cell immunity might be very long-lived. One report by Le Bert and colleagues shows that patients who recovered in 2003 from SARS-CoV-1 infection harbor SARS-CoV-2 cross-reactive T-cells 17 years later⁹⁵.
- Optimal T-cell epitopes vary by ethnicity and geography (MHC/HLA allele frequencies can be calculated using various tools⁹⁶. (MHC in humans is also called Human Leukocyte Antigen, or HLA). Epitopes providing extensive MHC coverage have been reported for many pathogens, including SARS-CoV-2 (for example, as published by Ferretti et al⁹⁷ and by Liu and colleagues⁹⁸). Such publications are useful for selection of T-cell epitope peptides for use in specific geographical regions and ethnicities. The publication by Ferretti and colleagues provides empirical validation of CD8 T-cell epitopes, broad coverage of the most common MHC haplotypes, and largely confirms the empirical findings of Le Bert and of

⁸⁹ <u>https://www.nature.com/articles/s41423-020-0401-3</u>

⁹⁰ https://www.frontiersin.org/articles/10.3389/fimmu.2020.00827/full

⁹¹ https://www.medrxiv.org/content/10.1101/2020.06.16.20130914v1

⁹² https://www.cell.com/immunity/fulltext/S1074-7613(20)30333-2

⁹³ https://www.medrxiv.org/content/10.1101/2020.05.08.20095430v3.full

⁹⁴ https://link.springer.com/content/pdf/10.1007/s12026-007-0071-6.pdf

⁹⁵ https://pubmed.ncbi.nlm.nih.gov/32668444/

⁹⁶ <u>http://allelefrequencies.net/</u>

⁹⁷ https://www.medrxiv.org/content/10.1101/2020.07.24.20161653v2

⁹⁸ https://www.cell.com/cell-systems/fulltext/S2405-4712(20)30238-6

Snyder et al⁹⁹. The study by Liu is computational and theoretically provides more complete haplotype coverage; however, only one of the SARS-CoV-1 cross-reactive T-cell epitopes described by Le Bert et al is present in this set of predicted epitopes, and it is predicted by Liu et al to be optimal for MHC1 and MHC2 haplotypes found mainly in Central Africa, a population not represented in the Singapore convalescents (mainly East Asian and European ancestry). And the top immunodominant MHC Class I / CD8 epitopes empirically verified by Snyder and colleagues are not represented in the computationally predicted set of Liu. Therefore, such computational predictions should be used with the caveat that they are not always consistent with empirical data.

- Peptide epitopes are preferred because they have the following advantages over other epitopes:
 - Well defined
 - Potentially pure
 - Enrichment of antigenic targets; not accompanied by substantial unnecessary and potentially counterproductive/non-neutralizing antigenic targets. Many epitopes are very short and well-represented by peptides (although uncertainty about epitope boundaries and addition of flanking amino acids to improve peptide solubility can warrant the selection of slightly longer sequences).
 - Some critical neutralizing B-cell epitopes can be very small, less than 20 amino acids.
 - The majority of CD4 T-cell epitope peptides are less than 20 amino acids
 - CD8 epitopes are often 9-11 aa.
 - Short production/turnaround time
 - \circ $\,$ Scalable; we currently use 50 to 100 micrograms total peptide per dose
 - Inexpensive relative to other epitope/antigen choices
 - Peptides allow specific mutations to be addressed in a modular fashion. The Gen 10+ vaccine designs include an epitope peptide with the increasingly common variant in the Spike RBD: N501Y. It is important to note that it is not necessary to include every mutated position of a variant of concern in order to neutralize the variant. For example, E484K is an increasingly common mutation in variants of concern, and appears to alter the interaction of Spike with the ACE2 receptor. Nevertheless, the widely studied neutralizing antibody REGN10987, and certain other neutralizing antibodies that bind the RBM, are not affected by mutations at position 484. Therefore, an epitope peptide that does not include position 484 of

⁹⁹ https://www.medrxiv.org/content/10.1101/2020.07.31.20165647v2

Back to top
the receptor binding motif, but does include the portion of the RBM bound by neutralizing antibodies should serve to elicit neutralizing antibodies, without regard to the identity of the amino acid at position 484.

- **Empirical evidence** should dominate selection criteria. Here are some best types of evidence:
 - Mapping of epitopes in blood and other samples collected from convalescent patients (ideally stratified by severity of illness). This can be accomplished by a few primary means:
 - 3D structural studies and modeling of neutralizing antibody binding to a viral antigen (e.g. Spike protein)
 - Mapping of linear B-cell epitopes by binding antibodies in convalescent sera to a library of peptides representing viral antigens. A strong signal in a linear epitope mapping study does not guarantee that the epitope peptide in the context of a vaccine will trigger the production of an antibody that binds to this epitope within the context of the virus. However, it is a good indicator that this is at least possible. Peptides can be constrained to approximate native conformation, making it more likely to bind the native epitope.
 - Mapping of T-cell epitopes by stimulating convalescent T-cells with epitope peptides, and measuring their response (e.g. cytokine secretion; ELISpot)
 - Epitope peptides from a peptide vaccine that has shown protection against infection
 - Successful use of epitope peptides in vaccines that elicit antibodies (or serum) effective in virus neutralization assays. B-cell epitopes that allow antibody binding to the virus but don't block viral function might increase risk of antibody-dependent enhancement.
 - Mapped epitopes that are effective in virus neutralization assays (e.g. peptides compete with viral sequences in cellular infection assays).
 - Successful use of epitope peptides in vaccines that elicit T-cell responses, or peptides shown to stimulate T-cells or cytokine production in ELISpot or other T-cell assay in cells from convalescents.
- Multiple epitopes with low mutant escape potential. It is important to note that most published neutralizing antibodies target Spike RBD, as do many vaccines in commercial development. However, given the high degree of mutability of the RBD portion of Spike, it is important to identify and select targets outside the RBD because of mutant escape potential. This strategy is summarized by Brouwer and colleagues: "In light of the rapid emergence of escape mutants in the RBD of SARS-CoV-1 and MERS, monoclonal NAbs

targeting other epitopes than the RBD are a valuable component of any therapeutic antibody cocktail. Indeed, therapeutic antibody cocktails with a variety of specificities have been used successfully against Ebola virus disease and are being tested widely in clinical trials for HIV-1. NAbs targeting non-RBD epitopes have been identified for SARS-CoV-1 and MERS, supporting the rationale..." for SARS-CoV-2¹⁰⁰. Three neutralizing epitopes described in the following section, containing or immediately adjacent to the Spike protease recognition sites and heptad repeat 2 (HR2) fusogenic domain are highly conserved among these 3 coronaviruses, suggesting that they have very low mutant escape potential.

- Two of these sites are cleavage sites for host cell proteases. Cleavage of the S1/S2 site by furin is required for subsequent cleavage of the s2' site by TMPRSS2. Cleavage by both is required for efficient fusion of the virus to the host cell membrane, and these sequences are constrained by recognition by these host proteases. Another region of the virus (heptad repeat 2 or HR2) is also required for membrane fusion, and the homologous 50 amino acid region of the SARS-CoV-1 virus is both 100% conserved, and is bound by neutralizing antibodies.
- According to 3D structural model data, relative to the RBD, these epitopes are structurally simple, and the peptides can be short (under 25 amino acids) and are water soluble. These features make them ideal for a peptide vaccine. The two protease sites loop out from the Spike protein surface, which makes them accessible to the required host proteases, but also to antibodies. The HR2 domain is predicted to form a coiled-coil "tether" that links the Spike trimer to the viral membrane. The linear epitope peptide should adopt a coiled coil conformation very similar to that exhibited within the native Spike protein.
- Therefore, rather than focusing on ACE2-binding epitopes in the highly mutation prone RBD to inhibit virus binding to the ACE2 receptor, we targeted these B-cell epitopes in the highly conserved portions of the Spike protein to strategically neutralize proteolytic cleavage and membrane fusion. Furthermore, all three are bound by antibodies present in the sera of large fractions of convalescents, and they produce among the highest signals in linear epitope mapping studies, which are far higher than signals measured for binding to any linear epitope in the RBD.
- The Gen 7 and 8 vaccine formulations still contain epitope peptides from within the RBD, including 2 peptides in the ACE2-binding region. However, these 2 peptides are unlikely to be conformationally correct and elicit antibodies that bind the native Spike conformation. The reason we include these is that they are

¹⁰⁰ <u>https://science.sciencemag.org/content/early/2020/06/15/science.abc5902</u>

Back to top

predicted to be effective CD4 T-cell epitopes, although empirical data from convalescent patients has not yet confirmed animal studies and computational predictions.

- **3D modeling of viral antigens** (especially Spike protein) is very helpful in B-cell epitope selection and peptide design.
 - Surface loop structures are often good antibody binding sites. Unstructured, hydrophilic loops that are stabilized by a minimal number of neighboring amino acid residues make good linear (rather than conformational) epitopes. One prime example in SARS-CoV-2 is Spike 804 to 818, the TMPRSS2 cleavage site (one of a few immunodominant epitopes reported by both Farrera et al and Li et al).
 - Certain amino acid residues are overrepresented and underrepresented in preferred antigens. According to structural studies, these amino acids are overrepresented (from most overrepresented): W, Q, P, K, S, E, N, H, C. And these are underrepresented (from most underrepresented): A, V, I, F, L, Y¹⁰¹.
 - There are multiple excellent 3D modeling packages and online servers. The NCBI supported iCn3d is very good and allows automatic access to Protein Database (PDB) files of viral protein structures¹⁰². Multiple 3D models of SARS-CoV-2 proteins are available, including Spike in various conformations, and complexed with antibodies and with the ACE2 receptor.
 - Conformational B-cell epitopes can be difficult to produce with peptides, but potentially might be designed as single peptides that join discontinuous portions of virus proteins, or can be produced as multiple peptides that are predicted to physically interact into relatively stable conformations.
- Physical properties of peptides. Certain peptides have excellent physical properties such as solubility and stability against proteolysis, and are synthesized easily; others are more problematic. Peptides with marginal properties can be improved by amino acid substitutions, additions, or subtractions. Bear in mind that changes to the viral sequence might substantially alter antigen recognition. It is likely preferable to include peptides that are in critical functional regions of the virus, with minor modifications to improve their use in a vaccine, than to exclude them. It is widely reported throughout the immunology and vaccine literature that both T and B-cell immunity are retained to varying degrees for new mutant strains. It has been reported that T-cell responses to SARS-CoV-2 are present in 34% to 60% of people not exposed to SARS-CoV-2, and are likely due to exposures to

¹⁰¹ <u>https://www.jimmunol.org/content/181/9/6230.long</u>

¹⁰² <u>https://www.ncbi.nlm.nih.gov/Structure/icn3d/full.html</u>

distantly related coronaviruses (most epitopes less than 50% aa identity)¹⁰³, ¹⁰⁴. Therefore, conservatively modified epitope peptides should be expected to provide high levels of protection, although possibly not equivalent to the unaltered viral amino acid sequences.

- Length. Binding of the S protein to the cellular receptor ACE2 is mediated through the receptor binding domain (RBD). The RBD has been mapped for the related SARS-CoV-1 virus, and for SARS-CoV-2. ACE2 binding residues in RBD have been published widely and can be visualized in Figure 3 of Tarek et al¹⁰⁵. However, multiple studies using short linear peptides for mapping epitopes bound by antibodies in convalescent sera reveal that only low signals are found in peptides within the RBD. This contrasts with several publications showing the isolation of neutralizing antibodies that bind to the RBD. Most of these mapping studies use epitope peptides of 12 to 15 amino acids, while Zhang et al use longer peptides of 20 to 25 amino acids and show that antibodies in convalescent sera show at least moderate binding to multiple RBD epitopes. Therefore, the peptides we have used in our vaccine formulations, especially in the RBD are longer than 20 amino acids, and as long as 40 amino acids.
- Solubility. Certain peptides of the high-scoring epitopes are predicted to have low solubility in water. This complicates formulation, since organic solvent or pH adjustment is required to solubilize. When choosing epitopes and peptides, bear in mind aqueous solubility. Solubility calculators are helpful in this process (for example, see ¹⁰⁶ and ¹⁰⁷). Marginal solubility of peptides can be improved by amino acid substitutions, additions, or subtractions. Acidic (Asp, Glu) and basic (Arg, His, Lys) amino acids aid solubility; combinations of acidic plus basic amino acids can keep the overall peptide charge near neutral.
- Presentation. B-cell epitopes presented on the surface of virus-like particles elicit stronger antibody responses than those encapsulated within particles. Both presentation and solubility can be enhanced by the synthesis of a terminal poly-acidic domain (e.g. several aspartic acid or glutamic acid residues), combined with a soluble chitosan derivative¹⁰⁸,¹⁰⁹.
- **Stabilization**. Circular peptides are more stable. Synthesis options are available to produce N-terminal to C-terminal cyclization, or to place cysteines in strategic

¹⁰³ <u>https://www.cell.com/cell/fulltext/S0092-8674(20)30610-3#articleInformation</u>

¹⁰⁴ <u>https://www.medrxiv.org/content/10.1101/2020.04.17.20061440v1</u>

¹⁰⁵ <u>https://www.medrxiv.org/content/10.1101/2020.04.25.20079426v1</u>

¹⁰⁶ <u>https://pepcalc.com/peptide-solubility-calculator.php</u>

¹⁰⁷ <u>https://www.thermofisher.com/us/en/home/life-science/[...]/peptide-analyzing-tool.html</u>

¹⁰⁸ <u>https://www.mdpi.com/2076-393X/3/3/730/htm</u>

¹⁰⁹ https://www.sciencedirect.com/science/article/abs/pii/S0968089619303347?

positions on epitope peptides for disulfide formation. Cyclization stabilizes 3D conformation, and increases resistance to exopeptidase degradation. Disulfide formation is quick, easy, and inexpensive by the treatment of peptides with activated charcoal. Cysteines should be designed into epitope peptides at positions that do not change or interfere with normal epitope function, or do so as little as possible. Replacement of amino acids with cysteine is indicated by the lower case letter "c" in certain epitope peptides. Blocking the N terminus of a peptide also has been reported to stabilize it against proteolysis. ¹¹⁰ Peptides might be either N-terminal acetylated or biotinylated.

We have produced several iterations of vaccine, and for each we selected multiple B-cell and T-cell epitopes. Which B-cell epitopes elicit antibody responses can be verified by immunoassay testing, and these epitope peptides can be carried forward in updated versions of vaccines. The use of multiple epitopes also ensures that, if vaccination is successful in eliciting immunity against, say, 2 or 3 epitopes, then a single mutation will not allow the virus to evade immunity. Mutating virus and epitope variants should be monitored (nextstrain.org, bioRxiv, etc), and new epitope peptides incorporated into evolving designs as needed.¹¹¹

Mapping of antibodies from people previously infected but recovered (convalescent) has shown that there are key regions for antibody binding. SARS and MERS are lethal coronaviruses that have been used for comparison and cross-reactivity of convalescent antibodies. Jiang et al used SARS-CoV-2 specific proteome microarrays to show that S and N proteins bind both IgG and IgM, and the C terminal portion of the N protein exhibits the highest convalescent serum antibody binding signal of all tested proteins.¹¹² Antibodies from convalescent patients have been mapped to short, linear epitope peptides by Poh and colleagues¹¹³ (S protein only), Wang et al ¹¹⁴, and the previously mentioned Zhang et al, Farrera et al, and Li et al.

Neutralizing (or protective) antibodies are preferred to guide B-cell epitope selection. Antibodies to certain functional regions of the virus will neutralize its activity and infectivity. Many antibodies bind to the virus and some will inhibit function but only a subset of all antibodies will truly neutralize the ability of the virus to infect cells and replicate. Ongoing research is helping to identify which portions of the virus are key to neutralizing activity. Neutralizing activity is not observed in antibodies associated with antibody-dependent enhancement of disease (ADE).

¹¹⁰ <u>https://www.pnas.org/content/pnas/102/2/413.full.pdf</u>

¹¹¹ <u>https://nextstrain.org/ncov</u>

¹¹² <u>https://www.medrxiv.org/content/10.1101/2020.03.20.20039495v1</u>

¹¹³ <u>https://www.biorxiv.org/content/10.1101/2020.03.30.015461v1</u>

¹¹⁴ <u>https://www.biorxiv.org/content/10.1101/2020.03.26.994756v1</u>

High-resolution linear epitope mapping by Wang et al was done with arrays of peptides covering the entire length of ten SARS-CoV-2 proteins. Length of peptides is 15aa, and peptides overlap by 5aa. Convalescent sera IgG and IgM were applied separately to the arrays. Binding events to a given peptide but not to adjacent peptides in the array indicate tight binding primarily to the central 5aa portion. Table 3 shows the 61 bound epitopes. Only five of these span more than one peptide. One in the S protein spans 15aa or three peptides (start = 806). These stretches are highly suggestive of a long linear epitope, whereas shorter epitopes are potentially a portion of a structural epitope. Moreover, the longer sequence provides a larger target. The longer S protein epitope binds both IgG and IgM, and a largely overlapping epitope binds convalescent antibodies in the study by Poh and colleagues, and produces the highest signal in the later and larger-scale linear epitope mapping study by Li and colleagues. The epitope at this position (about Spike 805-820) spans a key proteolytic cleavage site required for viral entry into host cells. We included this epitope in Gen 3+ vaccine formulas, and as updated and progressively rich scientific information has become publicly available, we have added additional B-cell and T-cell epitopes to successive generations of vaccine. Each of these additions is detailed with references in the sections below.

Peptides and vaccine composition (last updated 2022-01-14)

Peptide synthesis general guidelines

Peptides are the most expensive and only custom component of RaDVaC vaccine designs. They can be synthesized at various scales, from individual to population scale with relative ease. The price of a peptide varies considerably between suppliers, and depends on additional requirements such as purity and modification.

 Purification. Peptides do not need to be purified, although certain peptides will perform best if purified. In particular, longer peptides, B-cell epitopes, li-key peptides, and those predicted to be difficult to synthesize due to low coupling efficiency are ideally purified. Short peptides predicted to be easy to synthesize (generally due to minimal hydrophobic amino acids) do not require purification. Peptide "purification" is commonly misunderstood but simply means that full-length peptides are separated from partial peptides resulting from truncated synthesis. These truncation products are due to imperfect and sometimes highly variable amino acid coupling efficiencies during synthesis of the growing peptide chain. Since peptides will be chopped up by proteases, purification does not add much value unless predicted coupling efficiencies are low and the resulting fraction of full-length peptide is low. Estimated peptide synthesis coupling efficiencies can be predicted¹¹⁵. Again, for peptides predicted to be difficult to synthesize (especially B-cell epitopes and li-key peptides), consider purification.

- **Desalting**. Peptides should have TFA removal and replacement with acetate. This is commonly known as desalting or buffer exchange, and standard grade buffer exchange is relatively inexpensive.
- **Quality**. Peptides ideally should be GMP but high quality non-GMP peptides can be obtained. (GMP is typically quite expensive.)
- **Solubility**. Pay close attention to peptide solubility.¹¹⁶ Water solubility makes vaccine production substantially easier.

In summary, many peptides do not need to be purified, but they should have TFA removed and replaced with acetate, and should be water soluble. Water insoluble peptides require organic solvents, which complicate the overall vaccine formulation. Through experimentation with various peptides, and addition of soluble residues or substitution of hydrophobic residues with soluble ones, we are able to achieve solubility in water while retaining conservative positions of top scoring epitopes represented on these peptides.

Gen 10 and forward (updated 2022-01-14)

RaDVaC generations 1 to 9 vaccine designs consisted largely of incrementally improved epitope peptide selections, based on a transition from initial preprint and published computational predictions to a primary focus on empirical data. Gen 10+ designs provide significant improvements over previous generations in essential aspects of vaccine composition, beyond epitope selection. As with all aspects of the RaDVaC project, if you don't want to test all components of the present design, you are free to experiment with variations on the guidelines provided here. Our network of participating researchers appreciates any information you provide on your experimental variations, and Gens 10 and forward improvements are partly guided by information provided by ongoing experiments and data collection from the RaDVaC network.

The following general improvements have been made to the Gen 10 and forward vaccine designs, relative to previous designs:

• Improved solubility at physiological pH by the use of derivatized chitosan (HACC) instead of unmodified chitosan.

¹¹⁵https://www.thermofisher.com/us/en/home/life-science/protein-biology/peptides-proteins/custom-peptidesynthesis-services/peptide-analyzing-tool.html

¹¹⁶ <u>https://pepcalc.com/</u>

Back to top

- The use of li-key technology, to increase T helper activation, combined with reduced MHC Class II restriction to more robustly activate cytotoxic T lymphocytes and B-cells for antibody production.
- Surface display of antigens for improved antibody response.
- A smaller set of core peptides combined with a list of optional peptides, providing greater functionality and improved representation of common MHC Class I alleles.
- An optional epitope representing the Spike Receptor Binding Motif (RBM). The RaDVaC
 primary protective strategy remains focused on the more highly conserved epitopes
 involved in membrane fusion, but we are testing the potential of this epitope sequence to
 boost the systemic antibody response.
- Optional dendritic cell targeting peptides for delivering T-cell epitopes to dendritic cells, an important cell type in the presentation of T-cell antigens.

Vaccines prior to Gen 10 were designed simply to allow ease of reproducible production, and conservatively to minimize the likelihood of excess immunostimulation. Given results from hundreds of people who have taken these vaccines, we know they have been extremely safe (no serious adverse events reported), but the intranasally delivered vaccine is likely less effective than possible if antigen presentation had been more effective, and the T helper response had been stronger. We believe this to be the case since many people who have taken the nasal vaccine are testing negative for serum antibodies with commercial and lab ELISA tests, while many who inject the vaccine (subcutaneous or intramuscular) are testing positive (saliva testing appears to be providing evidence of mucosal response among a subset of researchers who have administered the vaccine intranasally). These antibody responses are important indicators that we will work to maximize, and consider together with other evidence of the effectiveness of RaDVaC vaccine designs, including the number of people taking vaccine, number of infections, and severity of infections. Encouraged by reports of parenteral response, we remain committed to improving the effectiveness of intranasal delivery, which is safer and likely to be more widely adopted. Therefore, the Gen 10 and forward designs implement multiple changes to address weak systemic antibody responses to nasal administration, but production of the vaccine remains relatively simple--albeit slightly more complex than previously, while maintaining the use of the same minimal set of equipment and materials.

As with Gen 9 and forward, the Gen 13 vaccine design consists of a group of core peptides, and a group of optional peptides. Many people desire to make vaccine but have limited resources, so we have made the core peptide group minimal (6 peptides), while retaining essential features and improving others. The optional peptide list provides a range of options for improved geographic and ethnic specificity. Omicron BA.5 has risen to predominance in much of the world.

Back to top

44

However, BA.2.75 is rising in frequency, and it might compete for dominance with BA.5 in some geographies. Therefore, the core peptide list has one key peptide, **RBMO4/5-sc**, specific for BA.5, but an optional peptide, **RBMO2-sc**, can be exchanged for **RBMO4/5-sc** in vaccines designed for use in those areas (or they can be used together). Some peptide sequences through Gen 9 are still used in the Gen 13 design, so for those who already have certain prior peptides, partial updating can be achieved by addition of peptides new to Gen 9 and forward--in particular, the li-key MHC Class II peptides described below. Most pre-Gen 10 peptides are usable in Gen 13 as optional or partially redundant additions. The Gen 9 and forward B-cell core epitopes are essentially the same but with additional sequences for improved display on the nanoparticle surface. Much published evidence shows that display of epitopes on the surface of a nanoparticle substantially enhances B-cell activation and antibody production^{117, 118}.

Those familiar with Gen 9 and prior vaccine designs will notice that sequences of B-cell epitopes at approximately Spike 805 and 1145 are shorter in Gen 10 and Gen 11. This is the result of a post-Gen 9 publication by Garrett and colleagues providing single amino acid resolution mapping of the boundaries of linear epitope binding¹¹⁹, and another by Ladner and colleagues providing amino acid resolution mapping across coronaviruses to discover the most cross-reactive conserved residues in these SARS-CoV-2 epitopes¹²⁰. Note that in these situations, flanking sequences might still be important for proper 3D conformation, but these data help define which amino acids are essential for antibody binding, and least likely to mutate away from recognition by neutralizing antibodies. These two B-cell epitopes at approximately Spike 805 and 1145 both bind neutralizing antibodies, and are much more highly conserved than other targets of neutralizing antibodies. Plus, antibodies targeting the HR2 domain and linker at 1145 neutralize a broader range of clinical isolates than antibodies targeting the S1 portion of Spike and the RBD¹²¹.

Immunogenic enhancements

To increase T helper activation in Gen 10 and forward, the clinically tested li-key technology has been implemented for a few CD4 and combination epitopes¹²². The li-key portion (the peptide LRMK joined to the N terminus by a 5 to 6 carbon linker) provides enhanced binding to MHC Class II molecules, and thereby increased CD4+ T helper activation¹²³. This promotes both CD8/CTL activation and antibody production from B-cell epitopes. Importantly, li-key has been

¹¹⁷ https://www.tandfonline.com/doi/full/10.1080/14760584.2019.1578216?

¹¹⁸ <u>https://www.pnas.org/content/pnas/109/4/1080.full.pdf</u>

¹¹⁹ <u>https://www.biorxiv.org/content/10.1101/2020.11.16.385278v1.full</u>

¹²⁰ <u>https://www.sciencedirect.com/science/article/pii/S2666379120302445</u>

¹²¹ <u>https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0050366</u>

¹²² https://www.tandfonline.com/doi/full/10.1517/14712598.6.12.1311

¹²³ https://pubmed.ncbi.nlm.nih.gov/10781856/

tested safely and successfully in many animal experiments, and in human clinical trials, and a SARS-CoV-2 vaccine design using li-key is currently being tested by a U.S.-China consortium¹²⁴. To further enhance antigen presentation and the T helper activation, we also provide sequences of optional dendritic cell targeting peptides (DCTP), which have been used in combination with li-key MHC to further boost its immunogenic effects.

The approach we describe for displaying antigens on the nanoparticle surface is an adaptation of two approaches: one that has been used in multiple published studies, and another that has shown promise in a published small-scale pilot study¹²⁵. The former approach involves creating peptides with long poly-anionic termini consisting of a stretch of glutamic or aspartic acid residues, and nanoparticles are formed by mixing the peptides with derivatized cationic chitosan absent another anionic crosslinker¹²⁶. The second approach, demonstrated in a study by Correia-Pinto et al., was designed to test the hypothesis that antigen display on the surface of nanoparticles increases antibody response to B-cell antigens, and their vehicle and antigen of choice of chitosan and synthetic peptides, respectively, provide a good model and protocol for emulation. The approach is quite simple: synthesizing multiple aspartic acid residues on the N terminus of epitope peptides, and then ionically binding the peptides into the surface of nanoparticles subsequent to particle formation. This publication is especially instructive since their chitosan nanoparticles were produced by sodium triphosphate ionic gelation. However, it is clear that their use of chitosan hydrochloride is not optimal, since chitosan hydrochloride is neutrally charged and water insoluble at physiological pH¹²⁷. Therefore, we have implemented two minor modifications of this approach using a water soluble chitosan derivative, and having the B-cell peptide participate in the ionic gelation formation of the nanoparticle (as described in a later section on vaccine preparation).

Readers of this publication will note that a Pan-DR Epitope peptide (PADRE) was utilized in this antigen surface presentation design. However, we have decided to use the li-key technology instead of the PADRE approach to enhance the T helper response for the following reason. Both PADRE and li-key appear to reduce MHC Class II binding affinity variability due to genetic variation. However, li-key allows the use of pathogen-derived epitopes, and potent MHC Class II epitopes of SARS-CoV-2 have been identified. Including these viral epitopes in a vaccine design instead of a non-viral epitope will provide better T helper response to infection by a closely

¹²⁴ <u>https://link.springer.com/article/10.1007/s10549-020-05638-x</u>

¹²⁵ https://www.mdpi.com/2076-393X/3/3/730/htm

¹²⁶ https://www.sciencedirect.com/science/article/pii/S174270611830566X

¹²⁷ <u>https://link.springer.com/article/10.1007/s002890050448</u>

related virus (e.g. SARS-CoV-2 mutants), since T memory cells will be cross-reactive to the coronavirus epitopes, but not to the PADRE epitope.

To keep the number of core peptides manageable, some of the epitopes with redundant or less common MHC restriction classes have been moved to the list of optional peptides. It is ideal to check the expected allele frequencies for MHC Class I for likely class restrictions of your ethnicity/geography, and to include the appropriate epitope peptides (see the Allele Frequency Net Database¹²⁸ and allele frequencies by geographic region¹²⁹). We also include the aforementioned optional dendritic cell targeting peptides (DCTP1 and DCTP2) that have been modified to attach to the surface of the chitosan nanoparticle.¹³⁰,¹³¹

Code key to the lists below:

- Amino acid numbering, sequences, and substitutions: Numbering of amino acids is based on the parental strain Wuhan-Hu-1. Due to deletions or insertions, numbering of specific amino acids can be different in certain variants of concern, but numbering of a given amino acid in this document will refer to the Wuhan parental strain numbering. For example L452R refers to a substitution of a leucine at position 452 of the parental strain, but this amino acid resides at position 449 of the delta variant of concern. Upper case amino acids letters in a sequence indicate wild-type sequence. Lower case letters in a peptide sequence indicate an amino acid substitution of the parental strain sequence.
- Linkers: The symbol {-} indicates a 3 to 6 carbon linker, and the symbol {6} indicates a 5 or 6 carbon linker.
- Surface display: The letter "s" at the end of a peptide name indicates surface display of an epitope peptide (by extension of the peptide with a poly-aspartic acid sequence). These poly-aspartic acid sequences are OPTIONAL but data suggests that surface display greatly enhances B-cell activation and antibody production.
- **Disulfide circularization:** The letter "c" at the end of a peptide name denotes circularization by disulfide linkage of a pair of cysteines (prior to Gen 10 these were indicated by the suffix "cir" and the # symbol; either order as a disulfide peptide or circularize with activated charcoal).
- **Purification:** Some of these peptides will perform substantially better if they are purified; in particular, most B-cell and li-key peptides will perform best if purified. For our most recently synthesized peptides, purification level is indicated. These purification levels are

¹²⁸ <u>http://www.allelefrequencies.net/</u>

¹²⁹ <u>http://allelefrequencies.net/top10freqs.asp</u>

¹³⁰ https://www.jimmunol.org/content/172/12/7425.long

¹³¹ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8210616/</u>

not provided as recommendations, only for data sharing. Note that of the OPTIONAL peptides, our most recent synthesis included a few of these, including the purified B-cell peptides **Spike 816-s** and **Spike N501Y-s**. We have not produced and tested all optional peptides; we only share these designs as prospects for future testing, or as improvements to previous designs.

- **li-key additions:** MHC Class II peptides have four amino acid li-key sequences (LRMK) attached to the N terminus by a 5 to 6 carbon linker, indicated by the {6} symbol.
- MHC Class I restriction: The binding preferences for MHC Class I and some MHC Class II
 peptides are shown for the most common alleles in most populations. Gen 11 Core
 peptides provide broad but not comprehensive coverage for many geographies and
 ethnicities. The optional MHC Class I peptides provide expanded ethnically specific
 coverage, while the optional combination MHC Class I/II li-key peptides also provide a
 wide selection of MHC classes and additional potential T helper cell activation on a single
 peptide.

Gen 13 updates (updated 2022-08-13)

The Gen 13 vaccine design is very similar to Gen 12, but with minor changes. Gen 12 was specific to omicron BA.1 and Gen 13 targets omicron BA.4-BA.5 (and/or, optionally, BA.2). The main change is the slight change of peptide **RBMO-sc**, which targets the receptor binding motif of BA.1, to **RBMO4/5-sc**. This peptide is predicted to be easier to produce by chemical synthesis. We are currently synthesizing this peptide. The 08-13 update provides an optional receptor binding motif peptide specific for BA.2.75.

Gen 13 Core peptides

- 1. **Spike 811-s**, dddddd{-}KPSgRSFIEDLLFNKVT-amide (combo B-cell/MHC Class I/II) (purified ≥85%)(MHC Class I: B40:01, B58:01)
- Spike 1146-s, acetyl-dddddd{-}DSFKEELDKYFKNHTSPD-amide (combo B-cell/MHC Class II) (purified ≥90%) (this peptide is slightly longer than its predecessor, Spike 1146-s, which can be used instead if you already have it)
- RBMO4/5-sc, dddddd { } cQSYGF PTyGVGhQP { 6 } NkLDSKVGGNYNY Yc-amide (B-cell, Receptor Binding Motif, Omicron; conformational epitope created by linking two discontinuous Spike peptides by the use of a 6 carbon linker. Residues shown in pink are omicron BA.4/BA.5 substitutions. Bold residues are ACE2 contact residues. (purified ≥90%) (MHC Class I sites: A01:01, A03:01, A11:01, A26:01, A30:02, A31:01, A32:01, A68:01, B15:01, B35:01, B57:01). Cysteine residues are included for circularization, but you may omit these if synthesis proves difficult.
 - <u>The similar RBMO-sc peptide failed chemical synthesis repeatedly. The reasons</u> for failure are very low coupling efficiencies, and side reactions of the poly-aspartic acid tail. Previous (non-omicron) versions of very similar RBM

peptides have displayed low coupling efficiencies but RBMO efficiencies were low enough that, coupled with this second problem of side reactions, synthesis was deemed too difficult to succeed without modification of the sequence. If you decide to synthesize this peptide as is, or with a modified sequence, please contact us at info@radvac.org.

- 4. **Mem 172T-key,** acetyl-LRMK{6}TSRTLSYYKLGASQRVA-amide (combo Class I/II; A11:01, A31:01, A30:01, A33:01, A33:03, B27:05, B39:01, C06:02, C07:01, C07:02) (purified ≥85%)
- Orf1ab 5528T DYGDAVVYRGTTTYKL (MHC Class I; A01:01, A03:01, A03:02, A11:01, A23:01, A24:02, A26:01, A30:01, A30:02, A32:01, A33:01, A33:03, A68:01, B15:01, B27:05, B35:01, B35:08, B39:01, B57:01, B58:01, C6:02, C07:01, C07:02)
- Orf1ab 4723T, RKIFVDGVPFVVSTGYHFRE (MHC Class I; A01:01, A02:01, A02:03, A02:06, A11:01, A23:01, A24:02, A26:01, A31:01, A32:01, A33:03, A68:01, B15:01, B35:01, B35:08, B53:01, C04:01, C05:01, C05:09)

Gen 13 Optional peptides

Targeting/effector peptides (typically test one or the other; we are testing DCTP2)

- DCTP1, acetyl-dddddd{-}rgFYPSYHSTPQRP-amide (dendritic cell targeting peptide 1) easy synthesis
- 2. **DCTP2**, acetyl-WPRFHSSVRHTH{-}ddddddd-amide (dendritic cell targeting peptide 2) easy synthesis

B-cell peptides

- 3. **Spike 816-s,** acetyl-SFIEDLLFNKVT{-}ddddddd-amide (B-cell, targeting the N terminal cleaved fusion peptide) possibly difficult synthesis (purified ≥85%)
- 4. **Spike 658-cs**, NSYECDIPIGAGICASYQT{6}QSaIAYT{-}ddddd (combo B-cell/CD4). Linker bypassing Q677 and the S1/S2 cleavage site.
- 5. DDDDDD{Ahx}CRSYSFRPTYGVGHQP{Ahx}NKLDSKVSGNYNYRYC
- 6. RBMO2.75-sc, dddddd { } cQSYGF PTyGVGhQP { 6 } NkLDSKVsGNYNYLYc-amide For areas with high levels of BA.2.75, this RBM B-cell epitope peptide is preferable to RBMO4/5, although the two can be combined into a single formulation. Mutations that differ from BA.4/BA.5 are shown in bright green. B-cell; conformational epitope created by linking two discontinuous Spike peptides by the use of a 6 carbon linker. Residues shown in pink are omicron BA.2 substitutions. Bold residues are ACE2 substituted contact residues. (purified ≥90%) (MHC Class I sites: A01:01, A03:01, A26:01, B15:01, B58:01, C6:02, C7:01, C7:02). Cysteine residues are included for circularization, but you may omit these if synthesis proves difficult.
 - <u>The highly similar RBMO-sc peptide failed chemical synthesis repeatedly. The</u> reasons for failure are very low coupling efficiencies, and side reactions of the poly-aspartic acid tail. Previous (non-omicron) versions of very similar RBM peptides have displayed low coupling efficiencies but RBMO efficiencies were low enough that, coupled with this second problem of side reactions, synthesis was

deemed too difficult to succeed without modification of the sequence. If you decide to synthesize this peptide as is, or with a modified sequence, please contact us at info@radvac.org.

Combination MHC Class I/II li-key peptides

- 7. **Nuc 102T-key,** acetyl-LRMK{6}KDLSPRWYFYYLGTGPEAGL-amide (A01:01, A23:01, A26:01, B07:02, B08:01, B15, B58:01), potentially difficult synthesis and purification
- Nuc 322T-key, acetyl-LRMK{6}MEVTPSGTWLTYTGAIKLDD-amide (A01:01, A26:01, B18:01, B35:01, B35:08, B40:01, B44:02, B44:03, B53:01), potentially difficult synthesis and purification
- Spike 1196-key, acetyl-LRMK{6}SLIDLQELGKYEQYIKWPWYI-amide (A01:01, A03:01, A03:02, A11:01, A23:01, A24:02, B15:01, B18:01, B44:02, B44:03, C06:02, C07:01, C07:02), potentially very difficult synthesis and purification

MHC Class I peptides

- 10. Orf1 5470T, KLSYGIATVR (MHC Class I; A02:01, A03:01, A03:02, A33:03)
- 11. Orf1 1636T, HTTDPSFLGRY (MHC Class I; A01:01, A03:02, A11:01, A26:01, A33:03)
- 12. **Orf8 107T,** DFLEYHDVRVVL (MHC Class I; A02:01, A23:01, A24:02, B39:01, B40:01, A33:03, C04:01, C05:01, C05:09, C06:02, C07:01, C07:02) Note: this region appears to be deleted in variant B.1.1.7 (UK); however, this deletion is associated with reduced clinical severity, so the inclusion of this peptide is worth considering.

The RaDVaC founding group is testing a Gen 11 vaccine including the core peptides plus **DCTP**, and also **Spike 816-s**, **MVC1-s** and **MVC2-s** for expanded antibody and mutational variant potential coverage, and **Nuc 264T-key**, **Nuc 102T-key**, **Orf1 5470T** and **Orf1ab 5528T** for expanded T-cell potential coverage due to the ethnic diversity of our group.

Spike or S protein amino acid sequences represented by epitope peptides

Numbered amino acid sequence for the S protein is shown below. (Sequences without breaks or spaces can be found in the Supplemental Materials section.)

Blue highlight shows the receptor binding domain (RBD), amino acids 319 to 529.¹³² Red highlight shows amino acids that contact ACE2, the receptor for the virus. Light yellow highlight shows the Heptad Repeat 2 (HR2) domain from aa 1163 to 1213. The S1/S2 protease cleavage site (rrar/sva) is shown in green highlight, with cleavage between Arg 685 and Ser 686. The S2' protease cleavage site (skr/sfi) is shown in magenta highlight, with cleavage between Arg 815 and Ser 816. The transmembrane domain starts at residue 1209, near the C terminus of the protein sequence. Underlined portions of sequence show coverage by epitope peptides.

¹³² <u>https://jvi.asm.org/content/jvi/early/2020/01/23/JVI.00127-20.full.pdf</u>

Wuhan parental strain

1	mfvflvllpl	vssqcvnltt	<u>rtqlppaytn</u>	<u>sftrgvyypd</u>	<u>k</u> vfrssvlhs
51	tqdlflpffs	nvtwfhaihv	sgtngtkrfd	npvlpfndgv	yfasteksni
101	irgwifgttl	dsktqslliv	nnatnvvikv	cefqfcndpf	lgvyyhknnk
151	swmesefrvy	ssannctfey	vsqpflmdle	gkqgnfknlr	efvfknidgy
201	fkiyskhtpi	nlvrdlpqgf	saleplvdlp	iginitrfqt	llalhrsylt
251	pgdsssgwta	gaaayyvgyl	qprtfllkyn	engtitdavd	caldplsetk
301	ctlksftvek	giyqtsnf <mark>rv</mark>	qptesivrfp	nitnlcpfge	vfnatrfasv
351	yawnrkrisn	cvadysvlyn	sasfstfkcy	gvsptklndl	cftnvyadsf
401	virgdevrqi	apgq <mark>t</mark> gkiad	ynyklpddft	gcviawns <mark>n</mark> n	ldskvggn <mark>y</mark> n
451	yl <mark>y</mark> r <mark>l</mark> frksn	lkpferdist	eiyqagstpc	ngveg <mark>fn</mark> c <mark>y</mark> f	pl <mark>q</mark> sygf <mark>q</mark> pt
501	<mark>ng</mark> vg <mark>y</mark> qpyrv	vvlsfellha	patvcgpkks	tnlvknkcvn	fnfngltgtg
551	vltesnkkfl	pfqqfgrdia	dttdavrdpq	tleilditpc	sfggvsvitp
601	gtntsnqvav	lyqdvnctev	pvaihadqlt	ptwrvystgs	nvfqtragcl
651	igaehvnns <u>v</u>	<u>ecdipigagi</u>	<u>casyqtqtns</u>	<u>prra</u> rsvasq	siiaytmslg
701	aensvaysnn	siaiptnfti	svtteilpvs	mtktsvdctm	yicgdstecs
751	nlllqygsfc	tqlnraltgi	aveqdkntqe	vfaqvkqiyk	tppikdfggf
801	n <u>fsqilpdps</u>	<u>kpskrsfi</u> ed	<u>llf</u> nkvtlad	agfikqygdc	lgdiaardli
851	caqkfngltv	lpplltdemi	aqytsallag	titsgwtfga	gaalqipfam
901	qmayrfngig	vtqnvlyenq	klianqfnsa	igkiqdslss	tasalgklqd
951	vvnqnaqaln	tlvkqlssnf	gaissvlndi	lsrldkveae	vqidrlitgr
1001	lqslqtyvto	q qliraaeira	a sanlaatkms	s ecvlgqskrv	/ dfcgkgyhlm
1051	sfpqsaphgv	v vflhvtyvpa	a qeknfttapa	a ichdgkahfp	o regvfvsngt
1101	l hwfvtqrnfy	y epqiittdnt	fvsgncdvvi	i givnntvydr	o lqpe <u>ldsfke</u>
1151	<u>eldkyfknht</u>	<u>sp</u> dvdlgdis	s ginasvvnic	q keidrlneva	a knlneslidl
1201	qelgkyeqyi	<mark>kwp</mark> wyiwlgf	f iagliaivmv	/ timlccmtso	c csclkgccsc
1251	gscckfdedd	d sepvlkgvkl	L hyt //		

Omicron variants^{133,134,135}



Omicron Spike mutations (relative to Wuhan-Hu-1) are shown in the above table for variants BA.1 through BA.5. BA.3 is a minor variant and is not shown in detail below.

BA.2



Spike mutations: T19I, L24S, Δ25-27, G142D, V213G, G339D, S371F, S373P, S375F, T376A, K417N, N440K, S477N, T478K, E484A, **Q493R, Q498R**, **N501Y**, **Y505H**, D614G, H655Y, N679K, P681H, N764K, D796Y, Q954H, N969K.

1	mfvflvllpl	vssqcvnl <mark>i</mark> t	<u>rtqsytn</u>	<u>sftrgvyypd</u>	<u>k</u> vfrssvlhs
51	tqdlflpffs	nvtwfhaihv	sgtngtkrfd	npvlpfndgv	yfasteksni
101	irgwifgttl	dsktqslliv	nnatnvvikv	cefqfcndpf	l <mark>d</mark> vyyhknnk
151	swmesefrvy	ssannctfey	vsqpflmdle	gkqgnfknlr	efvfknidgy
201	fkiyskhtpi	nl <mark>g</mark> rdlpqgf	saleplvdlp	iginitrfqt	llalhrsylt
251	pgdsssgwta	gaaayyvgyl	qprtfllkyn	engtitdavd	caldplsetk
301	ctlksftvek	giyqtsnf <mark>rv</mark>	qptesivrfp	nitnlcpf <mark>d</mark> e	vfnatrfasv
351	yawnrkrisn	cvadysvlyn	<mark>fa</mark> pf <mark>fa</mark> fkcy	gvsptklndl	cftnvyadsf
401	virg <mark>n</mark> ev <mark>s</mark> qi	apgq <mark>t</mark> g <mark>n</mark> iad	ynyklpddft	gcviawns <mark>nk</mark>	ldskvggn <mark>y</mark> n
451	yl <mark>y</mark> r <mark>l</mark> frksn	lkpferdist	eiyqag <mark>nk</mark> pc	ngv <mark>a</mark> g <mark>fn</mark> cyf	pl <mark>r</mark> sygf <mark>r</mark> pt
501	<mark>yg</mark> vg <mark>h</mark> qpyrv	vvlsfellha	patvcgpkks	tnlvknkcvn	fnfngltgtg
551	vltesnkkfl	pfqqfgrdia	dttdavrdpq	tleilditpc	sfggvsvitp
601	gtntsnqvav	lyq <mark>g</mark> vnctev	pvaihadqlt	ptwrvystgs	nvfqtragcl
651	igae <mark>y</mark> vnns <u>v</u>	<u>ecdipigagi</u>	<u>casyqtqt<mark>k</mark>s</u>	<mark>h</mark> rrarsvasq	siiaytmslg

¹³³ <u>https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/scientific-brief-omicron-variant.html</u>

¹³⁴ <u>https://www.ncbi.nlm.nih.gov/nuccore/OL672836</u>

¹³⁵ <u>https://outbreak.info/compare-lineages</u>

701	aensvaysnn	siaiptnfti	svtteilpvs	mtktsvdctm	yicgdstecs
751	nlllqygsfc	tql <mark>k</mark> raltgi	aveqdkntqe	vfaqvkqiyk	tppik <mark>y</mark> fggf
801	n <u>fsqilpdps</u>	<u>kpskrsfied</u>	<u>llf</u> nkvtlad	agfikqygdc	lgdiaardli
851	caqkfngltv	lpplltdemi	aqytsallag	titsgwtfga	gaalqipfam
901	qmayrfngig	vtqnvlyenq	klianqfnsa	igkiqdslss	tasalgklqd
951	vvn <mark>h</mark> naqaln	tlvkqlss <mark>k</mark> f	gaissvlndi	lsrldkveae	vqidrlitgr
1001	lqslqtyvtq	qliraaeira	a sanlaatkms	s ecvlgqskrv	v dfcgkgyhlm
1051	sfpqsaphgv	v vflhvtyvpa	a qeknfttapa	a ichdgkahfp	regvfvsngt
1101	hwfvtqrnfy	v epqiittdnt	fvsgncdvvi	givnntvydp) lqpe <u>ldsfke</u>
1151	<u>eldkyfknht</u>	<u>sp</u> dvdlgdis	ginasvvnic	y keidrlneva	a knlneslidl
1201	qelgkyeqyi	<mark>. kwp</mark> wyiwlgf	iagliaivmv	v timlccmtso	c csclkgccsc
1251	gscckfdedd	l sepvlkgvkl	hyt //		

BA.2.75¹³⁶,¹³⁷,¹³⁸

Spike mutations: T19I, L24S, Δ25-27, G142D, <mark>K147E</mark>, <mark>W152R</mark>, F157L, I210V</mark>, V213G, G257S, D339H, S371F, S373P, S375F, T376A, K417N, N440K, G446S, N460K, S477N, T478K, E484A, **R493Q**, Q498R, N501Y, Y505H, D614G, H655Y, N679K, P681H, N764K, D796Y, Q954H, N969K.

Pink highlights are omicron substitutions. Mutations above and below in bold type are substitutions of ACE2 binding residues. Mutations above in lime green highlight indicate amino acids mutated in BA.2.75 relative to BA.2. Note that R493 mutation in BA.2 has reverted back to the glutamine (Q) present in the Wuhan parental strain. Numbering is the same as the Wuhan strain, ignoring the altered numbering due to insertions and deletions.

1	mfvflvllpl	vssqcvnl <mark>i</mark> t	<u>rtq<mark>s</mark>ytn</u>	<u>sftrgvyypd</u>	<u>k</u> vfrssvlhs
51	tqdlflpffs	nvtwfhaihv	sgtngtkrfd	npvlpfndgv	yfasteksni
101	irgwifgttl	dsktqslliv	nnatnvvikv	cefqfcndpf	l <mark>d</mark> vyyh <mark>e</mark> nnk
151	s <mark>r</mark> mese <mark>l</mark> rvy	ssannctfey	vsqpflmdle	gkqgnfknlr	efvfknidgy
201	fkiyskhtp <mark>v</mark>	nl <mark>g</mark> rdlpqgf	saleplvdlp	iginitrfqt	llalhrsylt
251	pgdsss <mark>s</mark> wta	gaaayyvgyl	qprtfllkyn	engtitdavd	caldplsetk
301	ctlksftvek	giyqtsnf <mark>rv</mark>	qptesivrfp	nitnlcpf <mark>h</mark> e	vfnatrfasv
351	yawnrkrisn	cvadysvlyn	<mark>fa</mark> pf <mark>fa</mark> fkcy	gvsptklndl	cftnvyadsf
401	virg <mark>n</mark> ev <mark>s</mark> qi	apgq <mark>t</mark> g <mark>n</mark> iad	ynyklpddft	gcviawns <mark>nk</mark>	ldskv <mark>s</mark> gn <mark>y</mark> n
451	yl <mark>y</mark> r <mark>l</mark> frks <mark>k</mark>	lkpferdist	eiyqag <mark>nk</mark> pc	ngv <mark>a</mark> g <mark>fn</mark> cyf	pl <mark>q</mark> sygf <mark>r</mark> pt
501	<mark>yg</mark> vg <mark>h</mark> qpyrv	vvlsfellha	patvcgpkks	tnlvknkcvn	fnfngltgtg
551	vltesnkkfl	pfqqfgrdia	dttdavrdpq	tleilditpc	sfggvsvitp
601	gtntsnqvav	lyq <mark>g</mark> vnctev	pvaihadqlt	ptwrvystgs	nvfqtragcl

¹³⁶ <u>https://github.com/cov-lineages/pango-designation/issues/773</u>

¹³⁷ https://www.biorxiv.org/content/10.1101/2022.08.07.503115v1

¹³⁸ <u>https://www.who.int/activities/tracking-SARS-CoV-2-variants</u>

igae <mark>y</mark> vnns <u>v</u>	<u>ecdipigagi</u>	<u>casyqtqt<mark>k</mark>s</u>	<mark>h</mark> rrarsvasq	siiaytmslg
aensvaysnn	siaiptnfti	svtteilpvs	mtktsvdctm	yicgdstecs
nlllqygsfc	tql <mark>k</mark> raltgi	aveqdkntqe	vfaqvkqiyk	tppik <mark>y</mark> fggf
n <u>fsqilpdps</u>	<u>kpskrsfied</u>	<u>llf</u> nkvtlad	agfikqygdc	lgdiaardli
caqkfngltv	lpplltdemi	aqytsallag	titsgwtfga	gaalqipfam
qmayrfngig	vtqnvlyenq	klianqfnsa	igkiqdslss	tasalgklqd
vvn <mark>h</mark> naqaln	tlvkqlss <mark>k</mark> f	gaissvlndi	lsrldkveae	vqidrlitgr
lqslqtyvtq	qliraaeira	a sanlaatkms	s ecvlgqskrv	v dfcgkgyhlm
sfpqsaphgv	v vflhvtyvpa	a qeknfttapa	a ichdgkahfp	regvfvsngt
hwfvtqrnfy	v epqiittdnt	fvsgncdvvi	givnntvydp) lqpe <u>ldsfke</u>
<u>eldkyfknht</u>	<u>sp</u> dvdlgdis	ginasvvnic	q keidrlneva	a knlneslidl
qelgkyeqyi	<mark>. kwp</mark> wyiwlgf	iagliaivmv	v timlccmtsc	c csclkgccsc
gscckfdedd	l sepvlkgvkl	hyt //		
	igae <mark>y</mark> vnns <u>v</u> aensvaysnn nlllqygsfc n <u>fsqilpdps</u> caqkfngltv qmayrfngig vvn <mark>h</mark> naqaln lqslqtyvtq sfpqsaphgv hwfvtqrnfy <u>eldkyfknht</u> gscckfdedd	igae <mark>y</mark> vnns <u>y</u> <u>ecdipigagi</u> aensvaysnn siaiptnfti nlllqygsfc tqlkraltgi n <u>fsqilpdps</u> <u>kpskrsfied</u> caqkfngltv lpplltdemi qmayrfngig vtqnvlyenq vvnnnaqaln tlvkqlsskf lqslqtyvtq qliraaeira sfpqsaphgv vflhvtyvpa hwfvtqrnfy epqiittdnt <u>eldkyfknht</u> <u>sp</u> dvdlgdis gscckfdedd sepvlkgvkl	igae <mark>y</mark> vnns <u>v ecdipigagi casyqtqt</u> ks aensvaysnn siaiptnfti svtteilpvs nlllqygsfc tqlkraltgi aveqdkntqe n <u>fsqilpdps kpskrsfied llf</u> nkvtlad caqkfngltv lpplltdemi aqytsallag qmayrfngig vtqnvlyenq klianqfnsa vvnhnaqaln tlvkqlsskf gaissvlndi lqslqtyvtq qliraaeira sanlaatkms sfpqsaphgv vflhvtyvpa qeknfttapa hwfvtqrnfy epqiittdnt fvsgncdvvi <u>eldkyfknht sp</u> dvdlgdis ginasvvnic gscckfdedd sepvlkgvkl hyt //	igae <mark>y</mark> vnns <u>v ecdipigagi casyqtqtks hrra</u> rsvasq aensvaysnn siaiptnfti svtteilpvs mtktsvdctm nlllqygsfc tqlkraltgi aveqdkntqe vfaqvkqiyk n <u>fsqilpdps kpskrsfied llf</u> nkvtlad agfikqygdc caqkfngltv lpplltdemi aqytsallag titsgwtfga qmayrfngig vtqnvlyenq klianqfnsa igkiqdslss vvnhnaqaln tlvkqlsskf gaissvlndi lsrldkveae . lqslqtyvtq qliraaeira sanlaatkms ecvlgqskrv . sfpqsaphgv vflhvtyvpa qeknfttapa ichdgkahfp . hwfvtqrnfy epqiittdnt fvsgncdvvi givnntvydp . <u>eldkyfknht sp</u> dvdlgdis ginasvvniq keidrlneva . gscckfdedd sepvlkgvkl hyt //

BA.4, BA.5



The sequences of BA.4 and BA.5 Spike proteins are identical in most sequenced samples. Spike mutations: T19I, L24S, Δ25-27, Δ69-70, G142D, V213G, G339D, S371F, S373P, S375F, T376A, K417N, N440K, L452R, S477N, T478K, E484A, F486V, **Q498R**, **N501Y**, **Y505H**, D614G, H655Y, N679K, P681H, N764K, D796Y, Q954H, N969K.

Omicron BA.4 and BA.5 variants^{139,140}

Spike mutations: T19I, L24S, Δ25-27, Δ69-70, G142D, V213G, G339D, S371F, S373P, S375F, T376A, K417N, N440K, L452R, S477N, T478K, E484A, F486V, **Q498R**, **N501Y**, **Y505H**, D614G, H655Y, N679K, P681H, N764K, D796Y, Q954H, N969K.

Pink highlights are omicron substitutions. Mutations above and below in bold type are substitutions of ACE2 binding residues. Numbering is the same as the Wuhan strain, ignoring the altered numbering due to insertions and deletions.

1	mfvflvllpl	vssqcvnl <mark>i</mark> t	<u>rtq<mark>s</mark>ytn</u>	<u>sftrgvyypd</u>	<u>k</u> vfrssvlhs
51	tqdlflpffs	nvtwfhai	sgtngtkrfd	npvlpfndgv	yfasteksni
101	irgwifgttl	dsktqslliv	nnatnvvikv	cefqfcndpf	l <mark>d</mark> vyyhknnk
151	swmesefrvy	ssannctfey	vsqpflmdle	gkqgnfknlr	efvfknidgy
201	fkiyskhtpi	nl <mark>g</mark> rdlpqgf	saleplvdlp	iginitrfqt	llalhrsylt
251	pgdsssgwta	gaaayyvgyl	qprtfllkyn	engtitdavd	caldplsetk
301	ctlksftvek	giyqtsnf <mark>rv</mark>	qptesivrfp	nitnlcpf <mark>d</mark> e	vfnatrfasv
351	yawnrkrisn	cvadysvlyn	<mark>fap</mark> f <mark>fa</mark> fkcy	gvsptklndl	cftnvyadsf
401	virg <mark>n</mark> ev <mark>s</mark> qi	apgq <mark>t</mark> g <mark>n</mark> iad	ynyklpddft	gcviawns <mark>nk</mark>	ldskvggn <mark>y</mark> n
451	y <mark>ry</mark> r <mark>l</mark> frksn	lkpferdist	eiyqag <mark>nk</mark> pc	ngv <mark>a</mark> g <mark>vn</mark> c <mark>y</mark> f	pl <mark>q</mark> sygf <mark>R</mark> pt
501	<mark>Yg</mark> vg <mark>H</mark> qpyrv	vvlsfellha	patvcgpkks	tnlvknkcvn	fnfngltgtg
551	vltesnkkfl	pfqqfgrdia	dttdavrdpq	tleilditpc	sfggvsvitp
601	gtntsnqvav	lyq <mark>g</mark> vnctev	pvaihadqlt	ptwrvystgs	nvfqtragcl
651	igae <mark>y</mark> vnns <u>y</u>	<u>ecdipigagi</u>	<u>casyqtqt<mark>k</mark>s</u>	<mark>h</mark> rrarsvasq	siiaytmslg
701	aensvaysnn	siaiptnfti	svtteilpvs	mtktsvdctm	yicgdstecs
751	nlllqygsfc	tql <mark>k</mark> raltgi	aveqdkntqe	vfaqvkqiyk	tppik <mark>y</mark> fggf
801	n <u>fsqilpdps</u>	<u>kpskrsfi</u> ed	<u>llf</u> nkvtlad	agfikqygdc	lgdiaardli
851	caqkfngltv	lpplltdemi	aqytsallag	titsgwtfga	gaalqipfam
901	qmayrfngig	vtqnvlyenq	klianqfnsa	igkiqdslss	tasalgklqd
951	vvn <mark>h</mark> naqaln	tlvkqlss <mark>k</mark> f	gaissvlndi	lsrldkveae	vqidrlitgr
1001	lqslqtyvtq	qliraaeira	a sanlaatkms	s ecvlgqskrv	/ dfcgkgyhlm
1051	sfpqsaphgv	v vflhvtyvpa	a qeknfttapa	a ichdgkahfp	regvfvsngt
1101	hwfvtqrnfy	v epqiittdnt	fvsgncdvvi	i givnntvydp	o lqpe <u>ldsfke</u>
1151	<u>eldkyfknht</u>	<u>sp</u> dvdlgdis	s ginasvvnic	q keidrlneva	a knlneslidl
1201	. qelgkyeqyi	. kwpwyiwlgf	iagliaivmv	/ timlccmtso	c csclkgccsc
1251	gscckfdedd	l sepvlkgvkl	hyt //		

¹³⁹ <u>https://www.cdc.gov/coronavirus/2019-ncov/science/science-briefs/scientific-brief-omicron-variant.html</u>

¹⁴⁰ https://www.jpt.com/media/ce/9c/ca/1657637938/SpikeMutations_Overview.pdf

Nucleocapsid or N protein

Only T-cell epitope peptides were selected from the sequence of the Nucleocapsid protein. Underlined portions of sequence show coverage by epitope peptides. Omicron mutations: P13L, Δ 31-33, R203K, G204R. Numbering is the same as the Wuhan strain, ignoring the altered numbering due to deletion.

```
1 msdngpqnqr nalritfggp sdstgsnqng garskqr rpqglpnnta swftaltqhg
61 kedlkfprgq gvpintnssp ddqigyyra trrirggdgk mkdlsprwyf vylqtgpeag
121 lpygankdgi iwvategaln tpkdhigtrn pannaaivlq lpqgttlpkg fyaegsrggs
181 qassrsssrs rnssrnstpg sskrtsparm agnggdaala lllldrlnql eskmsgkgqq
241 qqqqtvtkks aaeaskkprq krt<u>atkaynv tqafgrrq</u>pe qtqgnfgdqe lirqgtdykh
301 wpqiaqfaps asaffgmsri gmevtpsgtw ltytgaikld dkdpnfkdqv illnkhid<u>ay</u>
361 <u>ktfpptepk</u>k dkkkkadetq alpqrqkkqq tvtllpaadl ddfskqlqqs mssadstqa
//
```

Spike region 439-507

<u>NNLDSKVGGNYNYLY</u>RLFRKSNLKPFERDISTEIYQAGSTPCNGVEGFNCYFPL<u>OSYGFOPTNGVGYOP</u> This region of the Spike protein comprises the majority of the Receptor Binding Motif, and contains many of the ACE2 receptor contact residues. This region comprises the "left shoulder", "neck", and "right shoulder" of the torso model of the Spike RBD (see Dejnirattisai et al.)¹⁴¹.

RBMO4/5-sc

ddddd{-}cQSYGF<mark>r</mark>PT<mark>y</mark>GVG<mark>h</mark>QP{6}N<mark>k</mark>LDSKVGGNYNY<mark>r</mark>Yc-amide <u>IN TESTING, vaccine Generation 13</u>

Residues shown in pink are omicron BA.4/BA.5 substitutions. Bold residues are ACE2 contact residues. (purified \geq 90%) (MHC Class I sites: Cysteine residues are included for circularization, but you may omit these if synthesis proves difficult.

This peptide represents the **R**eceptor **B**inding **M**otif of **O**micron. Similar to previous generation peptides such as N501Y-s, t is a conformational epitope created by linking two discontinuous Spike peptides by the use of a 6 carbon linker. The epitopes are in reverse order from their linear order in the Spike protein, with the N-terminal peptide start at 493, and the C-terminal peptide start at 439. This epitope represents the "right shoulder" and "neck" in the torso model of the RBD, which bind many neutralizing antibodies. This epitope contains the following omicron mutations: N440K, L452R, Q498R, N501Y, Y505H,

Therefore, we believe this minimal RBM epitope has the potential to induce potent neutralizing antibodies such as REGN 10987, and that the N501Y substitution will either not affect binding, allowing neutralization of virus with essentially any amino acid at position 501, or if Tyr 501 participates in antibody binding, it will enhance neutralization against the many emerging variants

¹⁴¹ <u>https://www.cell.com/cell/fulltext/S0092-8674(21)00221-X</u>

that carry this substitution. The N501Y substitution creates the peptide sequence QSYGFRPTY that is predicted to bind strongly to multiple allelic variants of MHC Class I receptors including A01:01, A03:01, A11:01, A26:01, A30:02, A31:01, A32:01, A68:01, B15:01, B35:01, B57:01.



Figure: Linker portion of the peptide Spike N501Y-s, shown in the context of the wild-type Spike protein (PDB ID 6XM4). In the peptide Spike N501Y-s, the C-terminus of Pro507 (shown here in red, pointing right) is covalently joined by a flexible carbon linker to the N terminus of Asn439 (shown in dark purple, pointing down).

The similar RBMO-sc peptide failed chemical synthesis. The reasons for failure are very low coupling efficiencies, and side reactions of the poly-aspartic acid tail. Previous (non-omicron) versions of very similar RBM peptides have displayed low coupling efficiencies but RBMO efficiencies were low enough that, coupled with this second problem of side reactions, synthesis was deemed too difficult to succeed without modification of the sequence. If you decide to synthesize this peptide as is, or with a modified sequence, please contact us at info@radvac.org.

RBMO2.75-sc, ddddd{-}c<mark>Q</mark>SYGF<mark>r</mark>PT<mark>y</mark>GVG<mark>h</mark>QP{6}N<mark>k</mark>LDSKV<mark>s</mark>GNYNYLYc-amide <u>OPTIONAL vaccine Generation 13</u> For areas with high levels of BA.2.75, this RBM B-cell epitope peptide is preferable to RBMO4/5, although the two can be combined into a single formulation. Mutations that differ from BA.4/BA.5 are shown in bright green. B-cell; conformational epitope created by linking two discontinuous Spike peptides by the use of a 6 carbon linker. Residues shown in pink are omicron BA.2 substitutions. Bold residues are ACE2 substituted contact residues. (purified \geq 90%) (MHC Class I sites: A01:01, A03:01, A26:01, B15:01, B58:01, C6:02, C7:01, C7:02). Cysteine residues are included for circularization, but you may omit these if synthesis proves difficult.

The highly similar RBMO-sc peptide failed chemical synthesis repeatedly. The reasons for failure are very low coupling efficiencies, and side reactions of the poly-aspartic acid tail. Previous (non-omicron) versions of very similar RBM peptides have displayed low coupling efficiencies but RBMO efficiencies were low enough that, coupled with this second problem of side reactions, synthesis was deemed too difficult to succeed without modification of the sequence. If you decide to synthesize this peptide as is, or with a modified sequence, please contact us at info@radvac.org.



Figure: Portion of the wild-type Spike Receptor Binding Motif (RBM) covered by REGN10987 (from PDB ID 6XM4). REGN10987 contact residues are N440, L441, K444, V445, G446, N448, Y449, Q498. Amino acid N501 is shown at upper left. The **Spike N501Y-s** peptide consists of the amino acids shown and should ideally assume this conformation, except for whatever alteration is due to the N501Y substitution; and, except for features not shown: 6 carbon linker covalently connecting P507 and N439; N501Y; poly-aspartic acid N-terminal segment; and C-terminal amidation. As can be seen, P507 and N439 are on the opposite side of the portion of the RBM from the amino acids contacted by REGN10987; thus, the linker should not perturb or disrupt immune recognition of this structure.

Spike region 658-695

658 NSYECDIPIGAGICASYQTQTNSPRRARSVASQSIIAYT

This epitope spans the S1/S2 site that is cleaved by the host furin protease, which is required for subsequent cleavage at the S2' site by the host TMPRSS2 protease (Spike region 805-830 epitope). It also includes a frequently observed variable amino acid: Q677. Omicron has 3 mutations in this region; therefore, we are omitting this region from the omicron-specific Gen 12 design. The peptide below has a chemical linker that spans the mutated portion containing the cleavage site.

Spike 658-cs

NSYECDIPIGAGICASYQT { - }QSaIAYT { - }ddddd <u>OPTIONAL vaccine Generation 10, 11, 12</u>

We have not yet synthesized or tested this peptide; we share this design as an update to the previous peptide **Spike 660**. **Spike 660** contains a common mutation at Q677 which is addressed in **Spike 658-cs** by the use of a linker that bypasses Q677 and the S1/S2 cleavage site, joining flanking sequences.

Spike region 802-830

802 FSQILPDPSKPSKRSFIEDLLFNKVTLADA

Contains the site of the S2' cleavage site of the viral fusion peptide by the host TMPRSS2 protease, which is essential for efficient membrane fusion and host cell infection. The S2' site is cleaved between Arg 815 and Ser 816 (SKR/SFI). An immunodominant epitope bound by convalescent antibodies reported separately by Farrera et al, Poh et al, and Wang et al. Poh and colleagues demonstrated that this epitope is bound by neutralizing antibodies¹⁴². A March, 2021 publication by Yang Li and colleagues showed that the peptide sequence from 815 to 829 (S2-23

¹⁴² <u>https://www.nature.com/articles/s41467-020-16638-2</u>

in their publication) IEDLLFNKVTLA does not elicit neutralizing antibodies in mice.¹⁴³ However, this peptide lacks key amino acids at the N terminus, and other groups have shown that a longer peptide is immunogenic. Therefore, it is important to include these additional N terminal amino acids. Predicted B-cell epitope by Grifoni et al. Includes a top B-cell epitope prediction from Lon¹⁴⁴ and from Khan¹⁴⁵. Peng and colleagues provide empirical data showing the sequence NFSQILPDPSKPSKR is a CD4 epitope.

Spike 811-s

ddddd{-}KPSgRSFIEDLLFNKVT-amide IN TESTING, vaccine Generation 10, 11, 12

Minimal epitope peptide containing the site of the S2' cleavage site of the viral fusion peptide, starting at Lys811. For optional enhanced display on the nanoparticle surface, this peptide also features an N-terminal poly-aspartic acid segment attached by a 3 to 6 carbon linker. Single amino acid resolution mapping data show that much of the loop between 805 and 817 is not necessary for typical antibody binding, and that R815 is the most N-terminal amino acid directly involved in antibody binding. Nevertheless, 3D models show that certain amino acids participate in stabilizing interactions with the antibody binding portion of the peptide. In particular, Lys811 forms a stabilizing salt bridge with Asp820. Lys 814 plays no role in the stabilization of the antibody binding region, and to prevent it from binding to the poly-aspartic N terminal portion of the peptide, it was replaced by glycine (K814G).

¹⁴³ <u>https://www.sciencedirect.com/science/article/pii/S2211124721002291</u>

¹⁴⁴ <u>https://www.biorxiv.org/content/10.1101/2020.04.03.022723v1</u>

¹⁴⁵ <u>https://www.biorxiv.org/content/10.1101/2020.05.03.074930v1</u>



Figure: Structure of Spike 811-827 (from PDB ID 6XM4). This structure shows the uncleaved S2' site and fusion peptide. Note the positions of K811 and R815, and their interactions with E819, D820, and F823. Cleavage of the S2' site separates this structure between R815 and S816, exposing the fusion peptide.

Spike 816-s

acetyl-SFIEDLLFNKVT{-}ddddddd-amide IN TESTING, vaccine Generation 10, 11, 12

This peptide is designed to present the N terminal end of the fusion peptide that results after TMPRSS2 protease cleavage of the S2' site. This structure mediates the initial stages of contact between the virus and the host cell membrane, leading to fusion of the viral and host cell membranes. The N terminal portion of this sequence is highly conserved among coronaviruses, forming a highly distinctive coil structure that penetrates the host cell membrane. Single amino acid resolution epitope mapping shows that this region contains key residues in the binding of neutralizing antibodies. Therefore, we hypothesize that the cleaved structure presents a uniquely

immunogenic structure that should be bound by antibodies with different affinities than those that recognize the uncleaved structure.



Figure: Structure of Spike 816-827 showing the N terminus of the S2' cleavage site and the fusion peptide (from PDB ID 6XM4). This represents the amino acids of the fusion peptide sequence in native conformation. Following cleavage the structure might undergo a substantial rearrangement upon separation from the rest of the Spike protein, resulting in a different structure from the one depicted here.

S 815-sc

acetyl-RSFcEDLLFNKVTLA{-}ddadddd IN TESTING, vaccine Generation 13

Epitope peptide containing the site of the S2' cleavage site of the viral fusion peptide, starting at Arg 815. For optional enhanced display on the nanoparticle surface, this peptide also features an N-terminal poly-aspartic acid segment attached by a 3 to 6 carbon linker. Single amino acid resolution mapping data show that much of the loop between 805 and 817 is not necessary for typical antibody binding, and that R815 is the most N-terminal amino acid directly involved in antibody binding.

SO 935-sc

cQDSLSSTASALGKLQDVVN<mark>H</mark>NAQALNT { - } ddaddadd

IN TESTING, vaccine Generation 13

Omicron variant (pink highlight represents omicron substitution Q954H) epitope peptide that forms a conformational epitope together with the fusion peptide (e.g. Spike 815-s).



Spike 1146-s

acetyl-ddddd{-}DSFKEELDKYFKNHTSPD-amide
<u>IN TESTING, vaccine Generation 12</u>

PREDECESSOR Spike 1147-s IN TESTING. vaccine Generation 10. 11

This peptide is slightly longer than its predecessor, Spike 1147-s, yet is shorter than the previous Spike 1145. We lengthened this peptide slightly because new data suggests antibodies might bind the additional C terminal residues, although several publications show binding to core residues present on Spike 1147-s (acetyl-dddddd{-}gSFKEELDKYFKNH-amide), which can be used instead if you already have it). Like other recent B-cell peptides, Spike 1146-s includes a

polyaspartic N terminus for enhanced nanoparticle surface presentation. Most 3D structural models do not include this portion of the Spike protein. However, PDB ID 6XR8 presents a clear model of this structure. This peptide forms an alpha helix, and the three alpha helices of the Spike trimer form a triple helix bundle. Note that each of the three helices is formed by an identical peptide sequence. The formation of this triplet structure is promoted by interactions of multiple aromatic and hydrophobic amino acid side chains in the interior, surrounded by aromatic amino acids and interhelix salt bridges on the outside of the bundle core. Single amino acid resolution mapping data show the boundary amino acids for antibody binding. Only the most highly conserved, transmembrane proximal two-thirds of the helix participates in binding. Therefore, we have omitted non-essential residues in this epitope peptide. Now includes 1155YFKNHTSP1162



Figure. Spike 1147-1159 from PDB ID 6XR8. This figure shows the triple helix bundle formed by this highly conserved portion of the Spike protein.

Spike 1145

1145 LDSFKEELDKYFKNHTSP IN TESTING. vaccine Generation 7. 8. 9

Immunodominant convalescent antibody-binding epitope described in linear epitope mapping studies by Farrera et al and Li et al. This region was not covered in the studies by Zhang et al. This epitope is immediately adjacent to 50 aa heptad repeat 2 (HR2) fusogenic domain, which is essential for fusion of coronavirus and host cell membranes, and is 100% conserved between SARS-CoV-1 and CoV-2 viruses¹⁴⁶. Data by Yi et al suggests that this epitope competes with neutralizing serum from convalescents, indicating that neutralizing antibody binds this sequence¹⁴⁷.

Nucleocapsid protein epitopes

Nucleocapsid is highly antigenic and produces large signals in ELISA of convalescent sera. Because of concerns about ADE, we are not using Nucleocapsid B-cell epitope peptides. If concerns about ADE risk are resolved, then these selections might be considered for use in a vaccine. However, as described below, certain portions of Nucleocapsid have been shown to be the longest-lasting T-cell epitopes, and we have selected these for eliciting durable T-cell responses.

Nucleocapsid proteins of SARS-CoV-1 and CoV-2 display antibody binding immunodominance, and the portions bound are highly similar, which isn't surprising given the high degree of conservation between these proteins¹⁴⁸. The portions of Nucleocapsid that produce high signal in epitope mapping studies (for CoV-1) are amino acids 147-170 and 362-412. CoV-2 mapping by Wang et al produced very similar high binding signals for the C-terminal portion¹⁴⁹. Mapping of CoV-1 Nucleocapsid protein antibody binding epitopes shows no binding in the region 100-122, and low binding in 321-345, which represent the longer T-cell epitopes included in Gen 7+ vaccine. A CD8 T-cell epitope (at CoV-2 N 359-369) at the border of the C-terminal antibody binding region was selected for inclusion in Gen 8+ vaccine designs.

T-cell epitope peptides

Mem 172T 172 TSRTLSYYKLGASQRVA OPTIONAL Generation 9

¹⁴⁶ <u>https://www.thelancet.com/journals/lancet/article/PIIS0140673604157887/fulltext</u>

¹⁴⁷ <u>https://www.tandfonline.com/doi/full/10.1080/22221751.2020.1815591</u>

¹⁴⁸ <u>https://jcm.asm.org/content/42/11/5309.full</u>

¹⁴⁹ <u>https://www.biorxiv.org/content/10.1101/2020.03.26.994756v1</u>

Mem 172T-key

acetyl-LRMK{6}TSRTLSYYKLGASQRVA-amide
IN TESTING, vaccine Generation 10, 11, 12

Combination MHC Class II/CD4 and MHC Class I/CD8 (A11:01, A33:03, B27:05, B39:01, C06:02, C07:01, C07:02).

The top ranking MHC Class II/CD4 epitope according to data published by Peng and colleagues, as well as the top-ranking MHC Class II/CD4 epitope among Adaptive Biotechnologies' data as of Sept 28, 2020. The peptide KEITVATSRTLSYYK and LSYYKLGASQRVAGD both are separately in the top 10 MHC Class II/CD4 epitopes identified by Tarke and colleagues in a whole proteome screen¹⁵⁰. These sequences clearly are among the most potent MHC Class II epitopes in the SARS-CoV-2 proteome. Inclusion of the li-key peptide (LRMK, attached to the N terminus by a 5 to 6 carbon linker) in a vaccine is predicted to substantially increase the MHC Class II/CD4 response.

Nuc 100-120

100 KMKDLSPRWYFYYLGTGPEAG IN TESTING, vaccine Generation 7, 8, 9

Nuc 102T-key

acetyl-LRMK{6}KDLSPRWYFYYLGTGPEAGL-amide

OPTIONAL, vaccine Generation 10, 11, 12

Combination MHC Class II/CD4 and MHC Class I/CD8 (A01:01, A23:01, A26:01, B07:02, B08:01, B15, B58:01)

One of the cross-reactive SARS-CoV-2 epitopes to T-cells from SARS-CoV-1 convalescent patients tested by Le Bert and colleagues¹⁵¹. T-cells from these patients are cross-reactive to homologous epitopes from SARS-CoV-2, which are highly conserved between the two viruses. Also identified by Peng and colleagues as an immunodominant CD4/MHC class II epitope and CD8/MHC class I epitopes. The sequence SPRWYFYYL was also identified experimentally by Ferretti et al as a CD8 epitope with predicted HLA restriction of B07. The second highest ranked CD8 epitope identified by Snyder et al. in their proteome-wide screen, although in subsequent larger public data releases from Adaptive Biotechnologies, its rank dropped to ninth¹⁵².

Nuc 263T

263 ATKAYNVTQAFGRRG

¹⁵⁰ <u>https://www.sciencedirect.com/science/article/pii/S266637912100015X</u>

¹⁵¹ <u>https://pubmed.ncbi.nlm.nih.gov/32668444/</u>

¹⁵² <u>https://www.medrxiv.org/content/10.1101/2020.07.31.20165647v2</u>

IN TESTING, vaccine Generation 8; OPTIONAL, Gen 9

Nuc 264T-key

acetyl-LRMK{6}ATKAYNVTQAFGRRG-amide **OPTIONAL**, vaccine Generation 10 Combination MHC Class II/CD4 and MHC Class I/CD8 (A23:01, A24:02, A33:03, B15:01, B35:01, B35:08, B53:01, C06:02, C07:01, C07:02) The full-length epitope ATKAYNVTQAFGRRG was predicted by Fast and Chen as a broad-coverage epitope for MHC-I (74%) and for MHC-II (100%). This sequence largely overlaps a top MHC Class II/CD4 epitope in the public dataset of Adaptive Biotechnologies (PRQKRTATKAYNVTQAFGR), and of Tarke et al. (KRTATKAYNVTQAFG), and one of the cross-reactive SARS-CoV-2 epitopes to T-cells from SARS convalescent patients tested by Le Bert et al (KAYNVTQAFGRRGPE). The sequence KAYNVTQAF is the third highest ranked CD8 epitope identified by Snyder et al, although in subsequent larger public data releases from Adaptive Biotechnologies, its rank dropped to nineteenth. Nuc 264T-key provides the li-key version of the same peptide sequence as Nuc 263T but fixes a previously incorrect starting position number. Inclusion of the li-key peptide (LRMK, attached to the N terminus by a 5 to 6 carbon linker) in a vaccine is predicted to substantially increase the MHC Class II/CD4 response. CAUTION should be exercised in the use of this peptide, since a similar sequence in SARS-CoV-1 has been implicated in suppression of cytokine signaling.¹⁵³

Nuc 321-345

321 GMEVTPSGTWLTYTGAIKLDDKDPN IN TESTING, vaccine Generation 7, 8, 9

Nuc 322T-key

acetyl-LRMK{6}MEVTPSGTWLTYTGAIKLDD-amide
OPTIONAL, vaccine Generation 10, 11, 12

Combination MHC Class II/CD4 and MHC Class I/CD8 (A01:01, A26:01, B18:01, B35:01, B35:08, B40:01, B44:02, B44:03, B53:01)

One of the cross-reactive SARS-CoV-2 epitopes to T-cells from SARS-CoV-1 convalescent patients tested by Le Bert and colleagues. T-cells from these patients are cross-reactive to homologous epitopes from SARS-CoV-2, which are highly conserved between the two viruses. This sequence contains a predicted dominant T-cell epitope of Grifoni et al (MEVTPSGTWL, HLA restriction B^{*}40:01) ¹⁵⁴. MEVTPSGTWL was also verified experimentally by Peng and colleagues as a CD4 epitope and CD8 epitope likely covering HLA type B^{*}40:01. MEVTPSGTWL is also among the top

¹⁵³ https://pubmed.ncbi.nlm.nih.gov/20976535/

¹⁵⁴ https://www.sciencedirect.com/science/article/pii/S1931312820301669

20 ranked CD8 epitopes identified by Snyder et al. Mateus et al identified the peptide PSGTWLTYTGAIKLD as a CD4 epitope, with common coronavirus cross-reactivity¹⁵⁵. Inclusion of the li-key peptide (LRMK, attached to the N terminus by a 5 to 6 carbon linker) in a vaccine is predicted to substantially increase the MHC Class II/CD4 response.

Spike 1196-key

acetyl-LRMK{6}SLIDLQELGKYEQYIKWPWYI{-}dddd
OPTIONAL Gen 11.12

Combination MHC Class II/CD4 and MHC Class I/CD8 (A01:01, A03:01, A03:02, A11:01, A23:01, A24:02, B15:01, B18:01, B44:02, B44:03, C06:02, C07:01, C07:02)

The sequence of this portion of Spike protein lies at the boundary of the viral exterior and includes a few amino acids of the highly conserved transmembrane domain of the protein. It is bound by antibodies in serum from a small fraction of convalescent patients. However, Adaptive Biotechnologies data indicate that it is the most important MHC Class I epitope on the Spike protein, strongly binding a large number of common allelotypes. Given the broad HLA/MHC Class I coverage provided by this peptide it will be an attractive addition to vaccine designs; however, it is predicted to be very difficult to synthesize. According to Class II data published by Adaptive and by Tarke et al, this sequence is only a minor contributor to the MHC Class II response; nevertheless, we have designed it as an optional li-key peptide because of its broad MHC Class I range.

Nuc 359T

359 AYKTFPPTEPK

IN TESTING, vaccine Generation 8. 9 OPTIONAL Gen 9

Class I MHC/HLA restriction: A03:01, A03:02, and A11:01

The sequences KTFPPTEPK / AYKTFPPTEPK are among the top ten ranked CD8 epitopes identified by Snyder et al. The sequence KTFPPTEPK was also identified experimentally by Ferretti et al as a CD8 epitope with predicted HLA restriction of A*03 and A*11. Also identified experimentally by Peng and colleagues as a CD8/MHC class I epitopes likely covering HLA types A*03:01 and A*11:01.

Orf1 1636T

1636 HTTDPSFLGRY

IN TESTING, vaccine Generation 8. 9, 10, 11, OPTIONAL: Gen 12

Class I MHC/HLA restriction: A01:01, A03:02, A11:01, A26:01, A33:03

The sequence HTTDPSFLGRY is the top-ranked MHC class I/CD8 epitope identified by Snyder et al in a proteome-wide screen from Adaptive Biotechnologies, although in subsequent larger

¹⁵⁵ <u>https://science.sciencemag.org/content/early/2020/08/04/science.abd3871</u>

public data release from Adaptive, its rank dropped to second. The sequence TTDPSFLGRY was also identified experimentally by Ferretti et al as a CD8 epitope with predicted HLA restriction of A*01. Peng and colleagues didn't include Orf1 in their analysis, explaining the absence of this epitope in their preprint. This epitope sequence is part of the polyprotein Orf1ab; the sub-protein and start position numbering are Nsp3, amino acid 818 (Nsp3 818).

Orf1ab 4723T

4723 RKIFVDGVPFVVSTGYHFRE

IN TESTING, vaccine Generation 10, 11, 12

Class I MHC/HLA restriction: A02:01, A11:01, A23:01, A26:01, A33:03, B15:01, B35:01, B35:08, B53:01, C04:01, C05:01, C05:09

This peptide sequence RKIFVDGVPFVVSTGYHFRE is among the top five MHC Class I/CD8 epitopes in Adaptive Biotechnologies' data. The core peptide FVDGVPFVV is water insoluble but the longer peptide is water soluble, and covers a wide range of common Class I allelotypes. This epitope is highly conserved and in a January 2022 publication Kundu et al. showed that FVDGVPFVV is cross reactive with common coronaviruses in people uninfected with SARS-Cov-2.¹⁵⁶

Orf1 5470T

5470 KLSYGIATVR

IN TESTING. vaccine Generation 9.10

Class I MHC/HLA restriction: A02:01, A03:01, A03:02, A33:03

This peptide sequence KLSYGIATV is among the top ten MHC Class I/CD8 epitopes in Adaptive Biotechnologies' data and covers Class I MHC/HLA restriction A*02 (KLSYGIATV). With the addition of the flanking C terminal Arg, it is water soluble, and also covers A*03 (KLSYGIATVR) and B*33. This epitope sequence is part of the polyprotein Orf1ab; the sub-protein and start position numbering are Nsp13, amino acid 146 (Nsp13 146).

Orf1ab 5528T

5528 DYGDAVVYRGTTTYKL

IN TESTING, vaccine Generation 12

OPTIONAL, vaccine Generation 10, 11

Class I MHC/HLA restriction: A01:01, A03:01, A03:02, A11:01, A23:01, A24:02, A26:01, A33:03, B15:01, B27:05, B35:01, B35:08, B39:01, C6:02, C07:01, C07:02

The peptide sequence DYGDAVVYRGTTTYKL is among the top 100 MHC Class I/CD8 epitopes in Adaptive Biotechnologies' data. It is highly conserved across coronaviruses.

¹⁵⁶ <u>https://www.nature.com/articles/s41467-021-27674-x#Sec17</u>

Orf8 107T

107 DFLEYHDVRVVL

IN TESTING, vaccine Generation 9; OPTIONAL, Gen 11, 12

Class I MHC/HLA restriction: A02:01, A23:01, A24:02, B39:01, B40:01, A33:03, C04:01, C05:01, C05:09, C06:02, C07:01, C07:02

This peptide sequence DFLEYHDVRVVL is among the top twenty MHC Class I/CD8 epitopes in Adaptive Biotechnologies' data. It is highly water soluble, and covers a wide range of common Class I allelotypes. **DELETED IN VARIANT OF CONCERN B.1.1.7**

Nsp7 21T

21 KLWAQCVQLH

IN TESTING, vaccine Generation 8; OPTIONAL, Gen 9

Class I MHC/HLA restriction: A02:01, B44:02, B44:03, A58:01

Placed into the list of optional peptides for Gen 9 due to absence from Adaptive Biotechnologies' MHC Class II/CD4 data release in late September 2020 (Orf1ab, which includes Nsp7 was not included in the generation of their data set). It is also absent from the MHC Class II data of Tarke et al., which does include Orf1ab and Nsp7¹⁵⁷ One of the cross-reactive SARS-CoV-2 epitopes to T-cells from SARS-CoV-1 convalescent patients tested by Le Bert and colleagues. The sequence KLWAQCVQL was also identified experimentally by Ferretti et al as the top-ranked CD8 epitope with predicted HLA restriction of A*02. The same sequence is among the top 20 ranked CD8 epitopes identified by Snyder et al.

Other possible epitopes and peptides, including high solubility synthetic peptides are listed at the end of this document in the Supplemental Materials section.

SUMMARY OF THE PROTECTIVE STRATEGY OF THE GEN 13 VACCINE DESIGN (updated 2022-07-20)

This section describes the evolution of the design strategy of the RaDVaC vaccine to achieve the goal of sterilizing immunity. The B-cell-based protective strategy of the first few generations of vaccine was focused on the Spike RBD interaction with the host ACE2 receptor. However, there are three points that make this strategy suboptimal.

 The ACE2 contact portion of Spike RBD (the receptor binding motif, RBM) has a low degree of conservation. Structural models show that CoV-1 contacts ACE2 slightly differently from CoV-2, and the related MERS virus interaction is even less conserved.

¹⁵⁷ <u>https://www.sciencedirect.com/science/article/pii/S266637912100015X</u>

These data suggest that this portion of Spike has higher mutant escape potential than more highly conserved portions of the virus.

- 2. Mutants with higher transmissibility and potentially higher pathogenicity have arisen in multiple amino acids in and around the RBM. Many mutations appear to alter the interaction of Spike with ACE2, and the binding of certain neutralizing antibodies.
- The full RBM is very difficult to produce as synthetic peptides. Even if produced, conformational correctness is a top concern with such a structurally complex portion of Spike, which is likely to be critical for eliciting an appropriate antibody response to this antigen.

Given these concerns, and the fact that early attempts at synthesis of RBD epitope peptides was expensive, inefficient, and complex, RBM peptides were retired in Gen 9. However, we continued to work on solving the problem of creating peptides to represent the RBM, and emerging variants of concern show that mutations in the RBD are important for transmission and pathogenicity. Therefore, for the Gen 10 and forward designs, we include a specially designed single peptide that represents key residues on the RBM. We use this simplified approach to represent a portion of the RBM shown to be bound by neutralizing antibodies, such as REGN 10987, while incorporating key mutations. There are many such mutations in this portion of the omicron RBM, and we have incorporated many in the omicron-specific RBM peptide RBMO4/5-sc. But this is not the only or even primary focus of the Gen 13 design. As with multiple prior generations, a major focus remains B-cell targets on the Spike protein known to be involved in fusion of the virus and host membranes; in particular, the two targets at about Spike 815 and 1145, with a lesser focus on the less conserved protease cleavage site at about 685. Here is a summary of key reasons for these foci:

- These portions of the Spike protein are highly to extremely highly conserved. The 50 amino acid Heptad Repeat 2 (HR2) domain is 100% conserved between CoV-1 and Cov-2, as is the immediately flanking sequence at 1145 to 1160. The S2' protease cleavage site is also highly conserved, and mutant escape potential is constrained because this site must be recognized and cleaved by the host protease TMPRSS2.
- 2. These 3 targets are required for membrane fusion. Even if one mutates to escape antibody binding, a remaining antibody bound site should inhibit efficient fusion.
- 3. The two primary target sequences form simple structures that are easily synthesized with relatively short peptides.
- 4. These 3 targets are strongly bound by antibodies in large fractions of convalescent patients, as described in the linear epitope mapping studies cited in the section above.
- 5. These 3 targets bind neutralizing antibodies, as described in the studies cited in the section above.

It is important to note a few points about this strategy. As linear peptides, Li et al. demonstrated that these individual epitopes do not produce strong neutralizing activity.¹⁵⁸ However, their fusion peptide epitope mapping created a division between two peptides directly in the center of the best peptide candidate. Their mapping of epitope SFKEELDKYFKNH did not have this problem. Nevertheless, they did not measure the combined neutralizing effects of these epitopes, which are expected to be synergistic, since they both target membrane fusion. However, it is being reported that the omicron variant of concern is largely resistant to current commercial vaccines and to previous infection, and omicron is heavily mutated in the N terminal domain and the RBD but not the C terminal domain or epitopes involved in fusion. Convalescents have a mix of antibody responses to fusion-related sights: about 45% have antibodies to the fusion peptide PSKRSFIEDLLF, and about 69% have antibodies to SFKEELDKYFKNH.

In addition to refocusing the B-cell targeting of the Gen 10 and Gen 11 vaccine designs, we maintain our focus on empirically validated T-cell epitopes. Similar to our strategy for B-cell epitope redundancy, we have included multiple T-cell epitopes for common alleles.

PREPARATION OF CHITOSAN NANOPARTICLE VACCINE (updated 2022-07-22; update includes note about the adjuvant imiquimod)

We have performed many iterations of vaccine preparation, with a range of ingredients and concentrations. The vaccine formula and production protocols below are simpler than earlier versions. Current versions use water soluble derivatized chitosan rather than unmodified chitosan, and this simplifies preparation since unmodified chitosan is only soluble at acidic pH.

Many published protocols exist for the robust production of nanoparticle vaccines from chitosan and its derivatives. Various parameters for creating nanoparticles of specific sizes and characteristics have been studied and optimized. Ideal target nanoparticle size is in the range of about 100 nm to 200 nm, which is in the size range of many viruses, including SARS-CoV-2, although larger nanoparticles have been reported to work well. Key parameters that influence particle size are chitosan concentration, chitosan to TPP ratio (charge ratio, or mass ratio, CS:TPP), salt concentration, and peptide or protein loading of the nanoparticle (higher load, larger size).¹⁵⁹ Chitosan nanoparticle formation is extremely robust to a wide range of ingredient concentrations.

Chitosan is a deacetylated form of chitin, which is found in mushrooms and the shells of crustaceans such as shrimp and crabs (seafood allergies are not allergies to chitin). Deacetylation

¹⁵⁸ <u>https://www.sciencedirect.com/science/article/pii/S2211124721002291</u>

¹⁵⁹ https://scholar.google.com/scholar?cluster=18336291994796435345&hl=en&as_sdt=0,22
produces free amino groups, and a high density of positive charges on chitosan. Protonation of the amino groups is pH dependent, and chitosan is generally soluble below pH 6, and insoluble at neutral or physiological pH (it has a pKa in the range of about 6.2 to 6.5). Therefore, many different chitosan derivatives have been developed (for review see Wang et al¹⁶⁰). Aside from pH/pKa differences, positively charged derivatives behave very similarly to unmodified chitosan; therefore, the following protocols sometimes use the general term chitosan which can include these higher solubility chitosan derivatives.

Many publications have described the production and properties of trimethyl chitosan (TMC), and quaternary or quaternized chitosan, including 2-hydroxypropyl-trimethyl ammonium chloride chitosans (HACC or HTCC). Unfortunately, suppliers of these derivatives are somewhat limited, but HACC appears to be more readily available than TMC. These derivatives provide superior performance relative to unmodified chitosan in vaccine production. To make a chitosan nanoparticle by the extremely common technique of ionic crosslinking, a solution of chitosan is mixed with negatively charged triphosphate (sodium triphosphate, STP a.k.a. tripolyphosphate, TPP) under rapid stirring or vortexing. Ionically bonded gel nanoparticles form spontaneously upon mixing with TPP, including when mixed together with peptide. The methods we use are adapted from published methods for creating peptide-loaded nanoparticles by ionic gelation by mixing a soluble chitosan derivative with poly-anionic peptides as crosslinker¹⁶³, ¹⁶⁴.

Because of its commercial availability and superior properties, we selected HACC for preparation of nanoparticles, which has been used widely in research, including for the production of vaccines that have shown excellent results in animal studies¹⁶⁵,¹⁶⁶. The acceptable properties of HACC span a wide range: molecular weight, approximately 100 kDa to 400 kDa; Deacetylation (DA) 70% to 90%; Degree of Substitution (DS) should be as high as possible, ideally >90% (the DS of the HACC we use is ≥90%, derived from Sigma-Aldrich product 417963, DA ≥75%). Therefore, we assume approximately 70% derivatized chitosan. Some publications report chitosan to TPP w/w ratios; however, degree of deacetylation can influence gelling behavior and nanoparticle size. Therefore, parameters influencing nanoparticle characteristics have also been reported using charge ratios. Chitosan is a polymer composed of D-Glucosamine monomer subunits, and the molecular weight of each monomer is 161.16 Da. The majority of chitosan literature is focused on

¹⁶⁰ https://www.mdpi.com/1422-0067/21/2/487/htm

¹⁶¹ <u>https://www.frontiersin.org/articles/10.3389/fmicb.2015.00372/full</u>

¹⁶² <u>https://www.ncbi.nlm.nih.gov/pubmed/?term=16380189</u>

¹⁶³ <u>https://www.sciencedirect.com/science/article/pii/S174270611830566X</u>

¹⁶⁴ <u>https://www.mdpi.com/2076-393X/3/3/730/htm</u>

¹⁶⁵ <u>https://www.sciencedirect.com/science/article/abs/pii/S0144861716304623</u>

¹⁶⁶ <u>https://www.sciencedirect.com/science/article/pii/S0165247819305863</u>

unmodified chitosan; therefore, we have been guided by the extensive literature on chitosan, plus recent work focused on HACC to guide our vaccine formulations using HACC.

HACC typically is produced by reacting chitosan with 2,3-Epoxypropyltrimethylammonium chloride (a.k.a. Glycidyl trimethyl ammonium chloride), which preserves the single positive charge and increases the molecular weight of each derivatized monomer by 151.63 g/mol. Therefore, for approximately 70% derivatized chitosan, the average mass per monomer is 267.3 g/mol, and the average mass per positive charge is 267.3 Da/0.7 = 381.9. TPP has a MW of 367.86 Da and a charge of -5. Therefore, the average mass per negative charge is 73.57. An equimolar charge ratio (positive to negative charge) is achieved at a CS:TPP mass ratio of approximately 5.2:1 (compared to the CS:TPP ratio for unmodified chitosan of about 2.8:1).

Nanoparticles are smallest when the chitosan is most dilute. An ideal range of chitosan concentrations is 0.5 mg/ml to 1.5 mg/ml. Given the molecular weight ratio of HACC:CS of about 1.8, it is unsurprising that published protocols for making HACC nanoparticles use higher concentrations. Given the range of key factors tested in previous publications, an ideal range for CS:TPP ratio (w/w) for 80% deacetylated chitosan is about 6:1 to 8:1. Hu et al report that a low CS:TPP ratio increases solution turbidity, probably by creating links between nanoparticles.¹⁶⁷ We have seen this phenomenon with a CS:TPP ratio of less than 4:1. Therefore a reasonable range of weight ratios for HACC:TPP should be approximately 10:1 to 14:1.

Chitosan to epitope peptide ratio (w/w). Ratios of chitosan:peptide have been reported in the literature from 1:1 to 25:1. The 1:1 ratio was achieved using chitosan/TPP as a carrier for insulin¹⁶⁸ and separately for ovalbumin. As the CS:peptide ratio approaches 1:1 loading capacity increases but loading efficiency decreases. Peptide loading of nanoparticles also increases the size of the nanoparticles. A 25:1 CS:peptide (w:w) results in approximately 50% increase in particle size, while a 10:1 ratio approximately doubles particle size. NaCl reduces particle size. About 20 to 50 mM NaCl should allow for small particle size, and substantial loading of peptide into the nanoparticle. However increasing NaCl concentration reduces zeta potential of nanoparticles, a measure of stability and resistance to aggregation. A ratio of 1:1 of chitosan to peptide was used by Nevagi and colleagues in self-assembling trimethyl chitosan nanoparticles that required no TPP or other additional ionic crosslinker¹⁶⁹. This was achieved by synthesizing epitope peptides with additional homopolymeric portions of acidic residues (5, 10, or 15 amino acid polyanions), which supplied sufficient anionic crosslinker for formation of TMC nanoparticles. Both poly-glutamic acid and poly-aspartic acid produced self-adjuvanting vaccine particles, with

¹⁶⁷ <u>https://scholar.google.com/scholar?cluster=10442461531761123020&hl=en&as_sdt=0.22</u>

¹⁶⁸ <u>https://pubmed.ncbi.nlm.nih.gov/10554100/</u>

¹⁶⁹ <u>https://www.sciencedirect.com/science/article/pii/S0968089619303347</u>

poly-aspartic acid producing the highest levels of IgG when used in a mouse model. Below, we provide a standard TPP crosslinking protocol for using standard synthetic peptides, and a protocol for hybrid TPP plus polyanion crosslinking using B-cell peptides with extended acidic homopolymers¹⁷⁰.

If the vaccine is being used in outbreak or pandemic conditions, to ensure that the vaccine is free of infectious virus contaminants during administration, solutions should be sterilized prior to mixing. Small volumes of chitosan/peptide and of TPP can be rendered free of infectious virus by heating in capped 15 ml conical tubes to 55° C for at least 15 minutes (longer with larger tubes), allowed to cool to room temperature, then mixed in a small sterile beaker while magnetic stirring rapidly (>500rpm) for a few minutes. Per Amidi et al, 1% w/v Tween 80 can be added to assist solubility for high concentrations of chitosan and/or peptide.

Storage of solutions

All solutions should be stored long term in a freezer at -20° C or below. Peptide and chitosan stock solutions can be divided into smaller aliquots to avoid repeated freeze/thaw cycles.

Stock solutions

Stock solutions only need to be prepared if you are planning to make large quantities of vaccine. If you are planning on making small-scale batches, concentrated stock solutions are probably unnecessary, and you can simply make working solutions. Nevertheless, accurately measuring the very small amounts of required materials is best done by making larger volumes. A vaccine recipe for about 10 ml (about 12 to 15 doses) is provided in a section below, and working solution recipes below allow you to make several of these batches. If you require much more than this, consider making stock solutions. Note that peptide stock and working solutions are the same.

Once prepared, heat stock solutions to 55° C (15 minutes for 50 ml conical tube; 10 minutes for 15 ml conical tube) to ensure that they are free of infectious virus. 5 M NaCl generally does not require heat sterilization but can be treated at the same time as an added safety precaution.

- 1) dH₂O
- 2) 5M NaCl (in dH₂O)
- 3) 10 mg/ml HACC in 100 mM NaCl; 50 ml HACC stock solution recipe (the HACC we have tested is soluble at this concentration but if your HACC is less soluble, try a lower concentration):
 - a) Add 49 ml dH_2O to a 50ml conical tube

¹⁷⁰ <u>https://www.mdpi.com/2076-393X/3/3/730/htm</u>

Back to top

- b) Add 1 ml 5M NaCl to the tube
- c) Weigh 500 mg HACC on a jeweler's scale, and add to the tube.
- d) Cap the tube and shake or vortex occasionally until the chitosan is dissolved. This probably will take at least 30 minutes. Heating helps speed the process.
- e) Dilute to desired working concentration. To make 50 ml working concentration of 2 mg/ml HACC in 20 mM NaCl, add 10 ml stock solution to a 50 ml conical tube, and then fill the tube to 50 ml with dH_2O
- 4) 5 mg/ml $Na_5P_3O_{10}$ (sodium triphosphate, a.k.a. tripolyphosphate, TPP) in dH_2O .
- 5) 1.0 mg/ml to 2.4 mg/ml peptide solution(s). Mix peptides into dH₂O. See protocol below for peptides that require disulfide circularization. A master mix of peptides can be made after each is suspended in water, or peptide powders can be combined into a single tube and resuspended together. For example, multiple peptide solutions of 1.5 mg/ml can be mixed together, and the final concentration of the peptide mixture will be 1.5 mg/ml. If you intend to make vaccine that utilizes T-cell peptides and B-cell epitope peptides in a different manner (for example, use of poly-anionic B-cell peptides as crosslinkers for enhanced surface presentation, as detailed in Protocol 2 below), then separate T-cell and B-cell peptide master mixes can be created. B-cell peptides with poly-acidic extensions (e.g. several sequential aspartate amino acids) are acidic. Some of these are marginally soluble in water (e.g. MVC1-s and MVC2-s) and solubility is aided by the addition of a small amount of NaHCO₃ (sodium bicarbonate, a.k.a. baking soda) to the peptide solution (try 5 mM).
 - a) Add sufficient water to a conical tube (leaving headspace for mixing)
 - b) Add peptide powder.
 - c) Cap and invert or shake the tube until all of the peptide is dissolved.
- OPTIONAL: solutions containing additional adjuvants. Some of the RaDVaC core team are currently producing and testing an Omicron BA.4/BA.5 vaccine that includes low doses of imiquimod.

Working solutions

Working solutions of HACC and TPP can be created from 5x stock solutions by adding 1 part stock solution to 4 parts water. For example, for 50 ml TPP working solution, add 10 ml TPP stock solution to 40 ml dH_2O . Individual working solution recipes are also provided in detail below.

- 7) HACC: 2 mg/ml HACC in 20 mM NaCl; 50 ml recipe:
 - a) To a 50 ml conical tube, add NaCl to 20 mM by one of the following methods:
 Pipette 200 microliters 5 M NaCl into the tube; or, weigh 59 mg NaCl and add to the tube

- b) Weigh 100 mg chitosan and add to the tube
- c) Add dH_2O to 50 ml (about 49.8 ml dH_2O)
- d) Cap and shake the tube vigorously for about a minute, and then occasionally until all of the chitosan is fully dissolved. This might take up to 30 minutes for larger clumps of chitosan to fully dissolve.

8) For recipe 1 below

- a) **Combined B-cell and T-cell epitope peptide mix:** 1.5 mg/ml peptide mixture (all peptides mixed together, e.g. equimolar or equal weight of each).
- b) Ionic crosslinker for recipe 1 below: 1 mg/ml TPP: recipe for 40 ml:
 - i) Add 40 ml dH_2O to a 50 ml conical tube
 - ii) Add 40 mg TPP to the tube
 - iii) Cap and shake the tube until all TPP is dissolved.

9) For recipe 2 below

- a) T-cell epitope peptide mix: 2 mg/ml peptide solution(s), prepared as above. All T-cell epitope peptides mixed together; it is easiest to use either equimolar or equal weight of each.
- b) TPP+PAP ionic crosslinker: 2:1 w:w ratio. Current testing range: 0.8 mg/ml to 1.0 mg/ml TPP, 0.4 mg to 0.5 mg/ml PAP: recipe for 10 ml:
 - i) Add 5.0 ml of 1 mg/ml B-cell epitope peptide (PAP) stock solution (either equimolar or equal weight of each peptide) to a 15 ml conical tube
 - ii) To the same tube, add 3.0 ml dH2O
 - iii) Add 2.0 ml of 5 mg/ml TPP stock solution
 - iv) Pipette to mix or cap the tube and gently invert to mix
- 10) OPTIONAL: 1M solution of sodium bicarbonate (NaHCO₃, baking soda); recipe for 10 ml:
 - a) To a 15 ml conical tube, add 10 ml dH_2O
 - b) Add 840 mg $NaHCO_3$ to the tube
 - c) Cap and shake the tube until all $NaHCO_3$ is dissolved.

Charcoal-assisted circularization of peptides

For peptides that require disulfide circularization. After circularization, charcoal can be filtered out although we find that freezing peptide solutions with charcoal does not appear to cause problems, and the charcoal remains intact in the bottom of the tube upon thawing.

- 1. Resuspend lyophilized peptide powder in dH_2O at a concentration of 1.5 mg/ml in a tube with headspace and a tight fitting cap.
- 2. Estimate the amount of peptide in the tube and weigh approximately 1 to 2 equal weights of small (average 0.5 to 1 mm diameter) granular charcoal (e.g. for 10 mg peptide, use 10

to 20 mg charcoal). If the charcoal granules are larger, they can be broken up to increase surface area. Add the charcoal to the peptide solution and tightly cap the tube.

 Place the tube on agitator (rocker, shaker, rotator, etc). Incubate at room temperature 3 to 5 hours, with mild agitation. Make sure that the charcoal is mixing through the peptide solution, and not simply sitting in the bottom of the tube. This can be done by placing the tube on its side and shaking, or by rocking or inverting the tube on a rotator.

Mixing the vaccine

Protocol 1: Ionic gelation with cationic HACC and anionic TPP as ionic crosslinker - sample protocol to make 10 ml of vaccine (about 12 to 15 doses plus nasal sprayer priming waste)

- 1. Heat the working solutions to 50 to 60 deg C (20 minutes for 50 ml conical tube; 15 minutes for 15 ml conical tube) to ensure they are virus free.
- 2. Sterilize the small beaker and stir bar by cleaning with soap and water, and then drying by wiping with an alcohol-soaked kimwipe or paper towel.
- 3. Place the magnetic stir bar inside the small beaker, and place the beaker onto a magnetic stir plate.
- 4. Add 8 ml of 1 mg/ml soluble chitosan (HACC) working solution to the beaker. Turn on the stir plate and slowly increase the stir speed to at least 500 rpm.
- 5. Add 1.0 ml peptide solution (total of 1500 micrograms peptide).
- 6. OPTIONAL adjuvant solutions should be added at this time.
- 7. Slowly add 1.0 ml TPP solution one drop at a time, and after the last drop is added continue to stir for at least 1 minute.
- 8. Test the pH of the vaccine. The pH should be in the range of 5.0 to 7.0. If necessary, adjust pH with white vinegar or $NaHCO_3$ solution. Only a very small amount should be required per 10 ml vaccine.
- Aliquot 800 microliters vaccine into sterile capped microfuge or 2 ml tubes. Only about 500 to 600 microliters of this will be used as vaccine; some will be lost due to sprayer priming and other waste.

Protocol 2: Ionic gelation with cationic HACC and poly-anionic peptides (PAP) plus TPP as ionic crosslinkers - sample protocol to make 14 ml of vaccine (about 25 doses plus nasal sprayer priming waste)

This protocol is used for improved surface presentation of B-cell epitopes, based on the aforementioned research by Nevagi and colleagues¹⁷¹, and Correia-Pinto et al¹⁷². It employs anionic homopolymeric stretches of amino acids at the termini of B-cell epitope peptides. T-cell epitopes do not need to be surface exposed on the nanoparticle for efficient processing and presentation. Therefore, T-cell peptides do not bear these polyanions and are mixed into the chitosan solution prior to ionic crosslinking, thereby encapsulating them in the nanoparticle. We have designed our B-cell epitope peptides to be synthesized with poly-anionic extensions of 6 to 8 amino acids, which are linked to the epitope sequence by a flexible 3 to 6 carbon linker. We named these poly-anionic peptides or PAPs. These PAPs including linkers can be produced by solid phase synthesis, and should not be more difficult or expensive than standard peptide synthesis.

To crosslink PAPs to the surface of the HACC nanoparticle, the PAPs are mixed together with TPP, and this mixture is added dropwise to the chitosan and T-cell peptide mixture to form gel nanoparticles. This results in encapsulation of the PAPs but also should result in binding of PAPs to the surface. Based on the work by Correia-Pinto and their use of a 10:1 or 20:1 w:w ratio of chitosan to PAP, we use this range for our initial experimentation. While their designs provide desirable features, the use of chitosan hydrochloride salt is incompatible with the strategy of retaining ionically bound surface epitopes at physiological pH. Therefore, our protocol is based on HACC, and its positive charge at physiological pH should retain surface-bound PAPs. During dropwise addition of TPP, nanoparticles begin to form at a chitosan: TPP ratio of about 15:1. Although nanoparticle formation for HACC hasn't been studied guite as extensively as for unmodified chitosan, it has been shown to be similar to that of unmodified chitosan, although differences in molecular weight to charge ratio and degree of substitution must be taken into consideration¹⁷³. Given these similarities and differences, we expect that nanoparticle formation by dropwise addition of TPP should begin at a HACC: TPP ratio of about 20:1. With the use of HACC, Correia-Pinto's protocol A (mixing the epitope together with TPP) should produce similar results to their protocol B (binding PAPs to the formed nanoparticle), because substantially over half of PAPs should remain in solution and be exposed to the surface after initial formation and ionic condensation of the nanoparticle. This should allow remaining PAPs to undergo equilibrium association with the nanoparticle surface. The range of HACC:TPP ratios we are testing has been

¹⁷¹ https://www.sciencedirect.com/science/article/pii/S0968089619303347

¹⁷² https://www.mdpi.com/2076-393X/3/3/730/htm

¹⁷³ <u>https://www.tandfonline.com/doi/abs/10.1080/10601321003742147</u>

Back to top

guided by prior publications on the formation of HACC nanoparticles by ionotropic gelation with TPP ¹⁷⁴, ¹⁷⁵, ¹⁷⁶.

Four ingredients are involved in nanoparticle formation, encapsulation, and ionic crosslinking:

- 1. HACC, 2-hydroxypropyltrimethyl ammonium chloride chitosans (also called HACC)
- 2. TPP
- 3. Crosslinking poly-anionic peptide (PAP) master mix of B-cell epitope peptides; about ¹/₃ of total peptide mass is poly-anionic
- 4. Non-crosslinking peptide (NCP) master mix of T-cell epitope peptides

HACC concentration: 2mg/ml w:w ratios: Ratio of HACC to TPP = In testing, a range of 10:1 to 12:1 Ratio of HACC to NCP = 5:1 Ratio of HACC to PAP = In testing, a range of 20:1 to 24:1 Ratio of TPP to PAP = 2:1

Premix a solution of 1 mg/ml each TPP and PAP.

- 1. Heat the working solutions to 50 to 60 deg C (20 minutes for 50 ml conical tube; 15 minutes for 15 ml conical tube) to ensure they are virus free.
- 2. Sterilize the small beaker and stir bar by cleaning with soap and water, and then drying by wiping with an alcohol-soaked kimwipe or paper towel.
- 3. Place the magnetic stir bar inside the small beaker, and place the beaker onto a magnetic stir plate.
- 4. Add 10 ml of 2 mg/ml HACC, 20 mM NaCl working solution to the beaker. Turn on the stir plate and slowly increase the stir speed to at least 500 rpm.
- 5. Add 2 ml 2.0 mg/ml T-cell peptide solution (total of 4.0 mg peptide) to the beaker.
- 6. OPTIONAL adjuvant solutions should be added at this time.
- 7. Add 2.0 ml PAP+TPP solution to the beaker. Slowly add solution one drop at a time, and after the last drop is added continue to stir for at least 5 minutes.
- 8. OPTIONAL: Test the pH of the vaccine. The pH should be in the range of 5.0 to 7.0. If pH is outside of this range, and if desired adjust with vinegar or NaHCO₃ solution.

¹⁷⁴ <u>https://www.mdpi.com/2073-4360/10/11/1226</u>

¹⁷⁵ <u>https://www.mdpi.com/1420-3049/26/2/449</u>

¹⁷⁶ <u>https://www.tandfonline.com/doi/full/10.3109/10717544.2013.853708</u>

 Aliquot 500 microliters vaccine into sterile capped microfuge or 2 ml tubes. Only about 400 microliters of this will be used as vaccine; some will be lost due to sprayer priming and other waste.

ADMINISTRATION OF THE VACCINE (updated 2021-03-07)

Dosage amount. Human studies have used as little as 7.5 micrograms inactivated virus or other replication incompetent antigen. Trials by El-Kamary et al. and Atmar et al. use 50 microgram and 100 microgram doses of viral-like particle in chitosan, although these trials use chitosan as a powder, which is a less efficient delivery vehicle than nanoparticles (see review by Smith et al.¹⁷⁷). However, the intranasal inactivated influenza vaccine trial in healthy human adults conducted by Illum and colleagues used 7.5 micrograms and 15 micrograms per dose, with greater than double vaccine effectiveness (VE) at the higher dose¹⁷⁸. We have used 50 micrograms to 100 micrograms total peptide for the initial priming and boost doses. Typical doses have been in the range of 70 micrograms. As of late March 2021, hundreds of self-experimenters have shown good tolerance for this dose. Assuming average coupling efficiencies for our synthetic epitope peptides of 70% to 80%, we expect the minimum peptide dose to be about 50 micrograms full-length epitopes. li-key peptide doses in Phase II clinical trials of cancer vaccines have used several doses of 500 micrograms per dose. Vaccine preparation protocol 2 allows for administration of about 200 micrograms per dose.

Booster schedule. The chitosan/peptide vaccine is only moderately immunogenic, and has proven to be weakly immunogenic for the stimulation of systemic antibody response. This contributes to short-term safety, but requires higher and more doses for efficacy than injected vaccines with potent adjuvants. The pre-Gen 10 vaccine has proven to be most effective if used at least 4 times (as shown by animal studies and human clinical trial data): a priming dose, and administration of 3+ booster doses. Six people who have taken RaDVaC vaccines and subsequently became infected have taken 3 or fewer doses, while many more who have taken four or more doses haven't reported an infection. Therefore, our standard regimen has been at least four doses. Beginning with Gen 10 we incorporated li-key technology, which should increase the immunogenicity of the vaccine, so fewer doses might suffice. People over the age of 60 or with complications that might attenuate immune response, might consider additional booster doses. Booster doses have been taken as soon as a few days after the prior dose, or as long as a

¹⁷⁷ https://www.tandfonline.com/doi/full/10.4161/hv.27449

¹⁷⁸ <u>https://www.sciencedirect.com/science/article/abs/pii/S0169409X01001715</u>

¹⁷⁹ <u>https://www.sciencedirect.com/science/article/pii/S0923753419356819</u>

¹⁸⁰ <u>https://link.springer.com/article/10.1007/s10549-020-05638-x</u>

few weeks. Ideally, doses are spaced by 1 week or more. Given the seriousness of the pandemic and individual circumstances, even shorter intervals between boosters might be used. This is done rarely even in a research setting, but most research is conducted in support of eventual commercialization, and reflects those constraints. Considering the burden and level of stimulation that a few weeks of CoV-2 infection exerts chronically on the immune system, more and frequent booster doses is not an unreasonable strategy. Nevertheless, there is some evidence of greater efficacy for longer intervals between doses. Dozens of RaDVaC testers have self-administered 4+ doses of progressive generations of vaccine, spaced by 1+ week each. Testing for immune response throughout the booster schedule and a few weeks after is ideal for monitoring building immunity; however, this is not practical for some people. Some have reported after-effects (e.g. pronounced nasal congestion) that increase with later booster doses. This is a positive qualitative sign of immune response, but even positive antibody and other test results should not be relied upon as correlates of protection (a perfect proxy for sterilizing immunity).

Pre-administration. To assess the pre-vaccination state of the immune system, one or more saliva samples, nasal wash or swab, and blood draws are ideal prior to vaccination.

Administration. Vaccination is achieved by nasal administration with the use of small (5ml to 20ml) commercial nasal sprayers. We have tried a few sources of spray bottles. Depression of the sprayer top of those we have tested delivers approximately 100 to 150 microliters of fine mist. This should be tested empirically by spraying into a small tube or beaker, and measuring the volume using a pipette. Dose is adjusted accordingly. Spray should be directed into each nostril. The sprayers we obtained come as a top sprayer unit, and either opaque white plastic, amber, or clear bottle. The sprayer unit stem stretches to the bottom of the bottle. The delivered volumes of vaccine will be too small to be delivered reliably using the bottle. Therefore, we do not use the bottle for vaccination, and instead place the sprayer stem directly into the smaller vaccine vial. The stem of our tested sprayer units are slightly longer than a 2 ml microcentrifuge tube. We pipette an aliquot of vaccine into the tube, then place the stem of the sprayer into the vaccine for spray administration.

Protocol for vaccine administration

To administer vaccine, it is ideal to wear gloves and have a spray bottle of 70% alcohol (ethanol or isopropanol) to sterilize your gloved hands, vials, and equipment. Select a well-lit work surface that can be sterilized with alcohol.

Required materials:

• Clean and well-lit work surface

- Gloves (nitrile, latex, etc)
- 70% alcohol (ethanol or isopropanol), preferably in a spray bottle
- Small (5ml to 20ml) commercial nasal sprayer bottle
- Vial of vaccine containing 1 dose plus extra for priming the sprayer (total, 800 microliter)
- Clean and preferably sterile paper towels or tissues
- OPTIONAL: saline wash solution and collection tube(s)
- OPTIONAL: rack or holder for vaccine vial
- Sterilize the work surface with alcohol. Place a paper towel or tissue on the clean work surface, and spray with alcohol. You can place items on the surface and use the towel or tissue to dab or wipe items (e.g. the sprayer stem) while maintaining sterility.
- 2. Wipe the vaccine vial with an alcohol wipe. Loosen the cap so that it can be removed by lifting it off, but leave the cap in place, and set the vial upright on the work surface. If you choose to use a tube holder or rack for the vial, make sure it is sterilized.
- 3. Blow your nose thoroughly with a tissue or paper towel to clear your nasal passages. Use an alcohol soaked wipe or tissue to clean the outer area of the nose and just inside each nostril. This will help prevent inadvertent contamination of the sprayer tip with infectious virus that might be present on or just inside the nose. We do not recommend cleaning deep into the nasal passage; this is likely to do more harm than good.
- 4. Sterilize your spray bottle by placing 2 to 4 ml of 70% alcohol into the spray bottle, screw on the cap, and deploy the sprayer until a fine mist is sprayed.
- Remove the spray top, and spray the residual alcohol. While you are doing this DO NOT TOUCH THE SPRAYER TIP OR STEM to maintain sterility. After sterilization or the following optional nasal wash, do not return the spray top to the bottle prior to using the vaccine.
- OPTIONAL. You can use the sterilized spray top apparatus at this point to spray 100 mM NaCl into nostrils for collection of nasal wash samples. Spray saline into nostrils, inhale slightly, and then evacuate nasal wash into a collection tube. Collect at least 500 microliters.
- 7. While holding the spray top in one hand, uncap the vaccine vial with the other hand, and place the clean and sterile stem of the sprayer directly into the vaccine vial so that the stem touches the bottom of the vial.
- 8. Grasp the vaccine vial with one hand, and the spray top with the other hand so that your index finger and middle finger are on the side handles of the sprayer.
- 9. Deploy the sprayer into the air (not up your nose) until you see a fine mist, in order to clear residual alcohol and/or saline from the sprayer.

- 10. Insert the spray tip into a nostril. While inhaling through the nose, depress the sprayer (ideally, a total of about 100 to 150 microliters per spray). Sniff in the vaccine. Repeat the dosing, for a total of 200 to 300 microliters into each nostril. The entire procedure can take seconds or minutes but don't let the vaccine drip out of the nose, and don't blow your nose for at least an hour or two.
- 11. **OPTIONAL**. In later booster doses you can collect nasal washes or saliva after vaccination to assess immune response. At 4 hours post vaccination, collect the sample as described in the previous OPTIONAL nasal wash. Cytokine levels in the sample can be measured as indicators of response, if you have the means to do this.

ASSESSMENTS OF IMMUNE RESPONSE (updated 2021-03-07)

There are multiple important immune responses to an effective vaccine, and, therefore, multiple measures of potential effectiveness. These are not measures of effectiveness, but measurable immune response correlates of protection.

- B-cell/Antibody response
 - Neutralizing vs non-neutralizing
 - Neutralizing titer
 - Avidity/affinity of antibodies
- T-cell response (often measured by cytokine secretion)
 - Cell mediated: Th1/Cytotoxic T Lymphocytes (CTL)/CD8+
 - Humoral: Th2/CD4+

Neutralizing antibodies provide protection against viral infection and replication. They bind to key functional parts of the virus. Non-neutralizing antibodies can be beneficial by marking a virus for destruction, but they do not interfere with virus activities such as infection and replication, and they are also associated with ADE. High antibody titer is generally required for optimal protection. Avidity or affinity of the antibody binding to an antigen is another important measure of an effective antibody. An ideal B-cell response involves high-avidity neutralizing antibodies that bind to multiple viral epitopes. The sections below address details of measuring each of these key metrics. T-cell responses are often measured by cytokine secretion, or by counting cytokine secreting cells.

Samples to be collected and tested

- Whole blood and serum
- Nasal wash
- Saliva

Baseline samples will be collected, and then post-vaccination samples will be taken at booster timepoints, and also at later dates. Each of the above sample types has been used successfully to detect viral RNA and antibodies.

Collection of samples for immune testing. Ideally, all samples will be collected immediately prior to vaccination, and at various time points after initial and booster vaccinations.

Blood is ideally collected by venipuncture and vacutainer, but can be collected by lancet (finger) and deposited into sterile microcentrifuge tubes, without preservative or clotting chemistry. Antibody assays such as ELISA, typically start at dilutions of 1:100 or 1:200 and thus can be run on a few microliters of serum. About 50% of whole blood is recoverable as serum. Therefore, a few drops of blood (30+ microliters) is sufficient for each individual at each time point, for each test. It is ideal to have enough serum for several tests. Blood and serum can be stored for relatively short periods (weeks) at -20°C until assayed. Whole blood or buffy coat are required for certain assays, such as ELISpot, that use peripheral blood mononuclear cells (PBMCs).

Nasal wash. Method adapted from de Fijter et al ¹⁸¹. Use nasal sprayer to deposit 200+ microliters of sterile 100mM saline into one nostril. Collect nasal wash effluent in a clean, labeled tube (a small funnel that fits into the collection tube is very helpful). Collection tube should be free of preservative, or contain non-denaturing preservative such as EDTA. Repeat the procedure with the other nostril. Repeat until you have collected a total of at least 0.5 ml. Don't collect more than 2 ml. Store at -20° C until further processed or assayed. Once all aliquots are collected, vortex with glass beads, then centrifuge the samples at 1000 g for 10 minutes at 4°C to clarify. Pipette the supernatants into fresh microcentrifuge tubes, and either refreeze at -20° C or process immediately.

Saliva. At least 0.5 ml should be deposited into sterile tubes, free of preservative, or containing a non-denaturing preservative such as EDTA. Store at -20° C until assayed. Depending on dilution factor (typically starting at 10 fold or greater), assays can be run on a few microliters of saliva.

Antibodies and B-cell immune response

Methods developed for assessing immunity to other respiratory viruses, such as influenza, can be adopted for use. Immunity will be assessed using standard indirect ELISA antibody assays (shown below) of serum, nasal swab or wash, and possibly saliva.

Variations of ELISA allow testing of the following:

¹⁸¹ <u>https://pubmed.ncbi.nlm.nih.gov/8872971</u>

Back to top

- Presence/absence of antibodies that bind a viral specific antigenic target
- Antibody titer (by dilution series)
- Antibody avidity (e.g. by pulse of chaotrope, such as thiocyanate or urea)

While neutralizing antibodies can't be identified by ELISA, this information can be inferred from published information on which epitopes are bound by antibodies that show neutralizing activity. Thus, if a specific epitope peptide used in a vaccine is bound by an antibody, we can infer that it should be a neutralizing antibody without directly testing it in a neutralization assay.

Initial ELISAs will be performed using recombinant Spike RBD as the target antigen. A few of our B-cell epitope peptides lie within the RBD, and this region is highly represented in immunodominant neutralizing antibodies¹⁸². Followup ELISAs will be performed on various antibody fractions with individual epitope peptides in 96 well plates. Serum will be diluted 1/200, and plated in a standard 2X dilution series. Nasal washes are performed by use of a nasal sprayer and 100 mM saline, and the samples are analyzed by ELISA essentially as by Barria et al.¹⁸³ For a review of assessing the effectiveness of nasal vaccines for influenza, see Gianchecchi et al¹⁸⁴.

¹⁸² <u>https://immunology.sciencemag.org/content/5/48/eabc8413</u>

¹⁸³ <u>https://academic.oup.com/jid/article/207/1/115/874878</u>

¹⁸⁴ <u>https://onlinelibrary.wiley.com/doi/full/10.1111/irv.12664</u>



Schematic of an indirect ELISA assay. Viral antigen (Ag) is attached to a surface (plate well). Patient serum is added to the plate well, washed and blocked. An enzyme-conjugated secondary antibody that binds to human antibodies is added to the plate well. A substrate for the enzyme is added to the plate well, producing a signal if the antigen is bound by antibodies in the patient's serum.

Standard protocols and reagents for testing for SARS-CoV-2 virus exposure have been published.¹⁸⁵,¹⁸⁶ These protocols and reagents allow testing for immunity either elicited from our vaccine or from virus exposure. Slight modification of the protocols to use our peptides in ELISAs, and for testing of nasal washes, will allow us to test for immunity specific to the epitopes of our vaccine, and for nasal mucosal immunity.

Antibody avidity can be measured by multiple methods, as reviewed by Klasse¹⁸⁷. In our vaccine schema, individual epitope peptides are used to elicit antibodies, and each of these can be used independently to assess avidity of binding antibodies.

B-cell epitopes and ELISA antigen targets

¹⁸⁵ <u>https://labs.icahn.mssm.edu/krammerlab/wp-content/[...]expression-and-serology-protocol-3 24 20.pdf</u>

¹⁸⁶ <u>https://labs.icahn.mssm.edu/krammerlab/covid-19/</u>

¹⁸⁷ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4766047/</u>

Spike protein epitopes / antigen targets

Purified recombinant Spike protein and Spike receptor binding domain (RBD) used as target antigen will allow for testing of viral exposure, and for immunity elicited by our vaccine, but will not be specific for immunity elicited by any specific epitope of our vaccine. Stadlbauer and colleagues use the recombinant RBD protein for testing rather than the full-length Spike protein, due to low expression and purification of full-length Spike. We will use RBD as a primary target antigen.

Epitope peptides used in vaccine production can be used in ELISA and other tests for immune response. ELISA testing with recombinant RBD and separately with epitope peptides might help determine if immunity is due to the vaccine, or to virus exposure. ELISA with a protein other than Spike will be more definitive in distinguishing between immunity from the vaccine, and immune response to virus exposure.

Nucleocapsid protein B-cell epitopes / antigen targets

Nucleocapsid is widely used as a highly sensitive antigen target for antibody testing. We ordered a peptide of one of the most antigenic sequences of the highly antigenic Nucleocapsid protein (NP produces large signals in ELISA of convalescent sera). Because of concerns about ADE, we have not yet included Nucleocapsid B-cell epitopes in our vaccine designs. It is possible to use them as negative controls for ELISA. If concerns about ADE risk are resolved, then these selections might be considered for use in a vaccine. We have designed longer peptides, and shorter portions of this peptide.

Nuc 371-399

Longer candidate: DKKKKADETQALPQRQKKQQTVTLPAAD

Long single linear epitope near the C terminus of the Nucleocapsid protein. Substitution of Lys 379 to reduce the positive charge near the N terminus: Lys>Gly. This region has the longest linear epitope discovered by protein microarrays in SARS-CoV-2 (Wang) and SARS-CoV (Zhu) ¹⁸⁸. This epitope is split into two overlapping peptides, **Nuc 376-399** and **Nuc 357-382. Nuc 376-399** has an additional Glutamate and Alanine at the N terminus to reduce the positive charge and help associate with chitosan.

Substituted Nuc 371-399: eaDKKKKADETQALPQRQKKQQTVTLPAAD

Nuc 357-382: IDAYKTFPPTEPKKDKKKKADETQAL

¹⁸⁸ https://www.pnas.org/content/103/11/4011.long

Prospective peptide for synthesis Overlaps with **Nuc 376-399**.

Nuc 371-394cir: Acetyl-cDKKKKADETQALPQRQKKQQTecL <u>SYNTHESIZED and ready for use as a negative control in testing</u> Nucleocapsid antibodies have been implicated in Antibody-directed enhancement. Take notice before selection for use in a vaccine. Useful as a negative control. Cys1, Cys24 disulfide, 2958 g/mol. 2.96g/ml = 1M. 2.96mg/ml = 1mM 1mg/ml = 338uM

Use of the recombinant RBD as a standard for ELISA will not allow high confidence that an immune response is due to the vaccine rather than virus exposure. Use of envelope or other non-Spike SARS-CoV-2 proteins in ELISA will be more definitive. A positive result in either or both Spike-RBD and Spike1, but negative for non-Spike proteins is suggestive of immune response due to the vaccine.

Positive controls for infection should be obtained. Blood and others samples should be obtained from people who are or have been infected. They should have systemic IgG antibodies to not only Spike protein, but to other proteins as well. They might also have sIgA antibodies in nasal wash and saliva.

T-cell immune response

There are many approaches to testing T-cell immunity specific to the virus. We are considering the following possibilities, and we welcome collaborations with scientists who can contribute to these and other analyses of T-cell responses.

- ELISpot¹⁸⁹
- Transcriptomic profiling of PBMC^{190,191}
- Immunome assessment

¹⁸⁹ https://en.wikipedia.org/wiki/ELISpot

¹⁹⁰ <u>https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3946932/</u>

¹⁹¹ <u>https://www.ncbi.nlm.nih.gov/pubmed/?term=30205979</u>

SUPPLEMENTAL MATERIALS

ADDITIONAL SEQUENCES

Wuhan strain Spike protein sequence

without breaks or spaces, for peptide search:

mfvflvllplvssqcvnlttrtqlppaytnsftrqvyypdkvfrssvlhstqdlflpffsnvtwfhaih vsqtnqtkrfdnpvlpfndqvyfasteksniirqwifqttldsktqsllivnnatnvvikvcefqfcnd pflqvyyhknnkswmesefrvyssannctfeyvsqpflmdleqkqqnfknlrefvfknidqyfkiyskh tpinlvrdlpqqfsaleplvdlpiqinitrfqtllalhrsyltpqdsssqwtaqaaayyvqylqprtfl lkynengtitdavdcaldplsetkctlksftvekgiygtsnfrvgptesivrfpnitnlcpfgevfnat rfasvyawnrkrisncvadysvlynsasfstfkcygvsptklndlcftnvyadsfvirgdevrgiapgg tqkiadynyklpddftqcviawnsnnldskvqqnynylyrlfrksnlkpferdisteiyqaqstpcnqv egfncyfplqsygfqptngvgyqpyrvvvlsfellhapatvcqpkkstnlvknkcvnfnfngltgtgvl tesnkkflpfqqfqrdiadttdavrdpqtleilditpcsfqqvsvitpqtntsnqvavlyqdvnctevp vaihadqltptwrvystqsnvfqtraqcliqaehvnnsyecdipiqaqicasyqtqtnsprrarsvasq siiaytmslgaensvaysnnsiaiptnftisvtteilpvsmtktsvdctmyicgdstecsnlllgygsf ctqlnraltqiaveqdkntqevfaqvkqiyktppikdfqqfnfsqilpdpskpskrsfiedllfnkvtl adagfikqyqdclqdiaardlicaqkfnqltvlpplltdemiaqytsallaqtitsqwtfqaqaalqip famqmayrfnqiqvtqnvlyenqklianqfnsaiqkiqdslsstasalqklqdvvnqnaqalntlvkql ssnfqaissvlndilsrldkveaevqidrlitqrlqslqtyvtqqliraaeirasanlaatkmsecvlq gskrvdfcgkgyhlmsfpgsaphgvvflhvtyvpageknfttapaichdgkahfpregvfvsngthwfv tqrnfyepqiittdntfvsqncdvviqivnntvydplqpeldsfkeeldkyfknhtspdvdlqdisqin asvvniqkeidrlnevaknlneslidlqelgkyeqyikwpwyiwlgfiagliaivmvtimlccmtsccs clkgccscgscckfdeddsepvlkgvklhyt//

Nucleocapsid protein sequence

without breaks or spaces, for peptide search:

msdngpqnqrnapritfggpsdstgsnqngersgarskqrrpqglpnntaswftaltqhgkedlkfprgqgvpintnss pddqigyyrratrrirggdgkmkdlsprwyfyylgtgpeaglpygankdgiiwvategalntpkdhigtrnpannaaiv lqlpqgttlpkgfyaegsrggsqassrsssrsnsstpgssrgtsparmagnggdaalalllldrlnqleskmsgk gqqqqgqtvtkksaaeaskkprqkrtatkaynvtqafgrrgpeqtqgnfgdqelirqgtdykhwpqiaqfapsasaffg msrigmevtpsgtwltytgaiklddkdpnfkdqvillnkhidayktfpptepkkdkkkkadetqalpqrqkkqqtvtll paadlddfskqlqqsmssadstqa